**United States Department of Agriculture**

Forest Service

North Central Forest Experiment Station

General Technical Report **NC-101**

# Data Bases For Forest Inventory in the North-Central Region

Jerold T. Hahn and Mark H. Hansen

LIBRARY FILE COPY
INTERMOUNTAIN STATION



North Central Forest Experiment Station

Forest Service--U.S. Department of Agriculture

1992 Folwell Avenue

St. Paul, Minnesota 55108

Manuscript approved for publication November 14, 1984

1985

CONTENTS


| | Page |
|---|---|
| I. Sampling and estimation procedures . . . . . | 3 |
| II. Codebook definition . . . . . | 4 |
| Format explanation . . . . . | 4 |
| Contents . . . . . | 5 |
| Record 1 (PLOTALL) . . . . . | 9 |
| Record 2 (PLOTFOR) . . . . . | 12 |
| Record 3 (TREE) . . . . . | 17 |
| Record 4 (SHRUBT) . . . . . | 20 |
| Record 5 (SHRUBL) . . . . . | 20 |
| Record 6 (SITETR) . . . . . | 21 |
| Record 7 (POINT) . . . . . | 22 |
| Index . . . . . | 23 |
| III. Stored procedures . . . . . | 26 |
| IV. Example retrievals . . . . . | 29 |
| Appendix . . . . . | 40 |
| A. Definitions of codes . . . . . | 40 |
| B. Glossary of terms . . . . . | 52 |
| Bibliography . . . . . | 57 |

# DATA BASES FOR FOREST INVENTORY IN THE NORTH CENTRAL REGION

Jerold T. Hahn, Principal Mensurationist,

and Mark H. Hansen, Mensurationist, Data Base Manager

The Forest Inventory and Analysis research work unit (FIA) at the North Central Forest Experiment Station, U.S. Department of Agriculture, Forest Service, collects information on various renewable natural resources in the 11-State North Central Region (fig. 1). To more efficiently use these data and to make them available to outside users, a commercially available data base management system was used to develop a set of data bases to enhance data storage and retrieval. This report is designed to be used by the managers, researchers, or planners who want to know what information is available from FIA data. Until recently, the only commonly available information was a series of reports that were published when a State




Figure 1.--Forest Survey units and data base boundaries.

Table 1.--Data base date, token, and size by geographic coverage

| | Geographic area covered | Date | Token[1/] | Bytes | Cases | Trees | Records | Forest plots |
|---|---|---|---|---|---|---|---|---|
| | | | | | | Number | | |
| I | Michigan Upper Peninsula | 1980 | MICHUP | 14,034,143 | 5,719 | 189,744 | 246,088 | 4,709 |
| II | Michigan Lower Peninsula | 1980 | MICHLP | 11,027,555 | 8,871 | 125,794 | 190,840 | 3,671 |
| III | Northern Wisconsin | 1983 | WISNOR | 15,060,621 | 7,151 | 200,351 | 274,679 | 5,425 |
| IV | Southern Wisconsin | 1983 | WISSOU | 6,576,303 | 8,768 | 89,651 | 135,884 | 2,790 |
| V | Northern Minnesota | 1977 | MINNOR | 17,014,071 | 13,696 | 235,885 | 274,804 | 8,345 |
| VI | Southern Minnesota | 1977 | MINSOU | 5,702,142 | 22,807 | 49,060 | 96,843 | 2,169 |
| VII | North Dakota | 1980 | NDAKOTA | 2,704,102 | 18,214 | 5,339 | 45,164 | 604 |
| VIII | South Dakota (eastern) | 1980 | SDAKOTA | 3,333,173 | 23,206 | 2,928 | 50,747 | 518 |
| IX | Nebraska | 1983 | NEBRASK | 2,314,016 | 14,449 | 5,004 | 35,906 | 535 |
| X | Kansas | 1981 | KANSAS | 5,371,730 | 23,705 | 36,306 | 93,931 | 1,807 |
| XI | Iowa | 1974 | IOWA | 2,562,666 | 12,607 | 16,276 | 43,303 | 756 |
| XII | Missouri Ozarks | 1972 | MOZARK | 4,323,264 | 4,042 | 67,956 | 78,148 | 2,108 |
| XIII | Missouri Riverborder and Prairies | 1972 | MORBNW | 2,403,360 | 6,569 | 27,145 | 41,470 | 884 |
| XIV | Illinois[2/] | 1962 | ILLINI | xx,xxx,xxx | x,xxx | xxx,xxx | xxx,xxx | x,xxx |
| XV | Indiana[3/] | 1967 | INDIANA | 1,500,725 | 1,058 | 24,124 | 28,028 | 1,058 |

[1/]Token used for data base name, password, and security passwords.
[2/]Data for Illinois are not expected to be available until 1985.
[3/]The Indiana data base contains data for forest plots only.

inventory was completed. Because of the extensive area covered by the inventories and the large variety of information gathered, only a small portion of the available information gets published. Persons needing different or more detailed data had to contact FIA personnel directly to see if the data were available and if so how they could obtain it. This report describes the FIA data available so that interested persons can decide whether or not those data can help satisfy their needs. In addition, it serves as a data base description for persons wishing to access the data bases directly. It is not meant to be a complete user's manual for the data base management system.

The FIA data bases are available through the University of Minnesota Computer Center (UCC) using the Scientific Information Retrieval (SIR) data base management system (Robinson et al. 1980). SIR is a hierarchical system designed for use with scientific data. Compared with other data base management systems available at UCC, SIR is the easiest, most economical, and most flexible system available for storing and retrieving FIA data. Interested users can directly access the FIA set of data bases and make their own data retrievals, or they can request the FIA data base manager to develop a specialized data retrieval for them. Users accessing the data bases will need to have a UCC user account, access to a computer terminal, knowledge of the SIR data base management system, and a basic understanding of FIA sampling and estimating procedures. The data base manager can assist anyone seeking a UCC account or information about SIR. Persons requesting the data base manager to access the data bases for them will be charged for computer and personnel costs.

The North Central FIA data base collection is stored in 15 separate SIR data bases, each covering a State or portion of a State (fig. 1, table 1). Within each data base is a data case for each inventory plot. Each case consists of up to seven data record types (table 2). These seven record types relate closely to the way data are collected and used in the FIA inventory.

This report is divided into four sections plus an appendix. The first section briefly describes FIA sampling and estimation procedures as they relate to the seven record types. The second section describes the data base, providing names, descriptions, formats, and data type codes for each variable in the data base.

Table 2.--North central FIA data base record types

| Record number | Record name | Record occurrence |
|---|---|---|
| 1 | PLOTALL | All ground plots |
| 2 | PLOTFOR | Forest plots and nonforest plots with trees plots |
| 3 | TREE | Measured trees on a plot |
| 4 | SHRUBT | Measured tall shrubs on a plot |
| 5 | SHRUBL | Measured low shrubs on a plot |
| 6 | SITETR | Site index trees recorded on a plot |
| 7 | POINT | Nonstocked points on a plot |

A table of contents and an index of variable names are included. The third section lists the stored procedures contained in the data bases and instructions on their use. The fourth section contains examples of retrievals that can be made from the data bases and demonstrates how interested users may access the data bases. The appendix describes those codes that vary between data bases or whose descriptions are too long to include in section II. The appendix also contains a glossary of terms and definitions used in this and other FIA publications.

## SECTION I.--SAMPLING AND ESTIMATION PROCEDURES

Users of the FIA data bases need to understand FIA sampling procedures in order to know the type of information that can be obtained from the data bases, how to go about obtaining this information, and the value and accuracy of the information. Specific FIA sampling procedures differ somewhat from State to State; however, the general description of the sampling procedure given here applies to all States.

The FIA inventory begins with an aerial-photo sample that estimates the area of land by various classes. These classes are based on land-use and, for forested land, forest type, volume, size and/or density. Also, as part of the aerial-photo sample, ground plots are located on the aerial photos and similarly classified. Ground plots may be remeasurement plots, transferred to the aerial photos from old photos or maps, or new plots, selected by systematic sampling. The photo classification of these ground plots, together with the area estimates from the photo sample, is used to assign area expansion factors to the ground plots. If 25 ground plots in the same sampling area have a photo interpretation classification that was observed for 50,000 acres in the sampling area, then each of these 25 ground plots would be given an expansion factor of 2,000 acres (50,000/25). The sampling area, or level at which expansion factors are assigned, is different from State to State, as is the classification scheme used to assign photo-interpretation classes. Sampling areas may be forest survey units (fig. 1), counties, groups of counties, or national forests. Persons interested in the details of how these expansion factors were assigned to the ground plots for a particular area should contact the data base manager.

Once a ground plot location has been selected on an aerial photograph, it is established and measured in the field. On all ground plots the information recorded includes plot location, land use, date of aerial photography, and date of ground measurement. This information is collected for all plots and is stored in record type 1 (PLOTALL), along with two expansion factiors. The first, the area expander (EXPAREA), is recorded for all plots and is used to estimate area based on all sample plots. The second, the remeasurement expander (EXPREMEA), is recorded only for remeasurement plots and is used to make estimates based on the remeasured sample.

On plots that are currently classified as forest or nonforest with trees, and on remeasurement plots that were forest at the previous measurement, additional plot information is recorded in record type 2 (PLOTFOR). This information includes ownership, total basal area, forest type, stand age, site index, and a volume expander (EXPVOL). The volume expander is similar to the area expander, and in most States the two are equal; however, in some States the inventory design necessitated a different expander. The volume expander must be used to expand plot volumes per acre to estimate total volume.

The basic FIA plot is a 10-point cluster that covers a 1-acre sample area. At each sample point a 37.5 BAF (English) prism plot is established, and all live trees with a diameter breast height (d.b.h.) 5 inches or greater are sampled, as well as salvagable dead trees or those that have died in the past 3 years. Trees with a d.b.h.less than 5 inches but at least 1 inch are sampled on fixed-radius (6.8 feet) plots. Stumps cut in the past 3 years are recorded on ten 1/50-acre plots. Each tallied tree or stump is identified by a unique point and tree number. Items tallied include species d.b.h., crown ratio, and other tree information. This information is stored in record type 3 (TREE). There is one occurrence of this record for each tree. Some of the items in this record, such as tree volume, are computed from basic tree measurements. There is also an expander for each tree (EXPANDR) that expresses the number of trees per acre represented by the sample tree.

Shrubs are sampled on the northeast quarter of a 6.8-foot fixed-radius plot located at each of the first three cluster sample points. Shrubs are tallied as tall or low. Tall shrub species and stem diameter are recorded in record type 4 (SHRUBT), and low shrub species and percent cover are recorded in record type 5 (SHRUBL).

Also, for each forest plot a number of additional trees are measured to determine site index. This information is recorded in record type 6 (SITETR) and occurs once for each site index tree that was measured. In addition, when one of the 10-point cluster sample points is observed to be nonstocked (no trees or seedlings present), an explanation of why it is nonstocked is recorded in record type 7 (POINT).

For remeasurement plots additional information is recorded for each tree, such as original tree d.b.h. and other original tree data. This information may be used to estimate growth, mortality, and removals on these remeasurement plots. On new plots removals can be estimated using the measured stumps and assuming these

stumps were cut in the past 3 years. Mortality on new plots can be estimated from the dead trees tallied, also assuming these trees died in the past 3 years.

Another method to estimate mortality using new plots is by using the mortality factor, an estimate of the number of trees per acre represented by the tally tree that will die in the next year. This mortality factor is the product of the tree expander and the probability that the tree will die in the next year. Growth on new plots is estimated using data stored as original tree data. For new plots, the original tree data are an estimate of the trees' condition 1 year before being measured, based on growth functions. These growth functions and the functions used to compute probability of mortality are derived from remeasurement plots. Details about these functions for a particular data base can be obtained by contacting the data base manager.

Finally, the data bases also contain projected plot and tree data that are periodically updated using the Stand and Tree Evaluation and Modeling System (STEMS) (Belcher et al. 1982) growth model. The system used to update the data simulates natural growth and mortality but does not account for removals, land-use change, regeneration, fire, management, or any other major changes that could occur. Methods are being developed to incorporate these major changes into the update process.

Not all data were collected for all surveys. For example, shrub data were first collected during the 1980 Michigan survey and are only available for this and subsequent surveys. If a data item is not available for all surveys, this is noted in the codebook (section II).

## SECTION II.--CODEBOOK DEFINITION

### Codebook Format Explanation

The codebook contains detailed definitions of the variables in the data base. A detailed explanation of this codebook can be found in the SIR user's manual (Robinson et al. 1980). We present a brief explanation of this format using the definition of sample kind as an example.

```
R2.      SAMPKND , Sample Kind
                 FORMAT:  I1
              DATA TYPE:  I*1
           VALUE LABELS:

                (1) New 10-Point Full
                (2) Remeasurement
                (3) Remeasurement NF
                (4) New 10-Point Full NF
                (5) Reserve [Phantom]
                (6) Remeasurement Partial
                (7) New 10-Point Partial
                (8) Remeasurement Partial
                (9) Remeasurement Updated
                (0) Stereo dots [not gr.ck]
```

Each definition starts with a variable number consisting of a letter (C or R) and one or more digits. A description of the difference between C (common information record) and R (record) variables can be found in Robinson et al. 1980. In the example the variable number is R2. Next is the variable name, one to eight characters all upper case, SAMPKND in the example. The name is followed by the variable label. This label can be of any length and consists of two parts: A short label of up to 40 characters followed by a semi-colon and as many 80-character lines separated by semi-colons as desired. The example has a short label only. This short label is the label transferred with the variable during retrievals. The next line shows the data input format for the variable, in this case a one-digit integer number, followed by the data type and data base storage requirement, in the example an integer variable using one byte of storage. Finally, the value labels are listed. These labels are up to 20 characters long and are listed for each legal value. In the example, we have labels for each of the 10 possible values. In general, if a value label does not exist, it means that the remaining values are illegal for that variable.

NORTH CENTRAL FIA DATA BASE CODEBOOK DEFINITION

TABLE OF CONTENTS

RECORD 1 (PLOTALL)


| Variable name | Variable type & number[1/] | Definition | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 9 |
| STATE | R1. | State Code | 9 |
| UNTCO | C1. | Unit and County | 9 |
| SAMPKND | R2. | Sample Kind | 9 |
| DISTURB | R3. | Disturbance Code | 9 |
| PICODE | R4. | Photo Interpretation Code | 9 |
| PISTRAT | R5. | Photo Stratification Code | 10 |
| GLUCUR | C2. | Ground Land Use Current | 10 |
| GLUORG | R6. | Ground Land Use Original | 10 |
| GLUPRJ | R7. | Ground Land Use Projected | 10 |
| CHANGE | R8. | Reason for GLU change | 10 |
| UTREND | R9. | Use Trend | 10 |
| STDHIST | R10. | Stand History | 10 |
| PRJSTDH | R11. | Projected Stand History | 10 |
| PHOTAGE | R12. | Photo Age | 10 |
| SURDATE | R13. | Survey Date | 10 |
| PRJDATE | R14. | Projected Date | 10 |
| GROPER | R15. | Remeasurement Period | 10 |
| USESET | R16. | Use Setting-Recreation | 10 |
| USEAREA | R17. | Use Area-Recreation | 11 |
| USEPOST | R18. | Use Posted-Recreation | 11 |
| SOILCLS | R19. | Soil Classification | 11 |
| EXPAREA | C7. | Area Expander | 11 |
| PRJAREA1 | R20. | Projected Area Expander 1 | 11 |
| PRJAREA2 | R21. | Projected Area Expander 2 | 11 |
| UTMZONE | R22. | UTM Zone Code | 11 |
| UTMNORTH | R23. | UTM Northing To Nearest 1000 Meters | 11 |
| UTMEAST | R24. | UTM Easting To Nearest 1000 Meters | 11 |
| TOWNSHIP | R25. | Township | 11 |
| RANGE | R26. | Range | 11 |
| SECTION | R27. | Section | 11 |
| EXPREMEA | R28. | Remeasurement Expander | 11 |


RECORD 2 (PLOTFOR)


| Variable name | Variable type & number | Definition | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 12 |
| DWT | R1. | Distance to Water Type | 12 |
| DWS | R2. | Distance to Water Size of Water | 12 |
| DWV | R3. | Distance to Water Value | 12 |
| DRT | R4. | Distance to Road Type of Road | 12 |
| DRV | R5. | Distance to Road Value | 12 |
| NATFOR | R6. | National Forest Code | 12 |
| RANDIST | R7. | Ranger District | 12 |
| OWNGP | C3. | Ownership Group | 12 |
| PRJOWN | R8. | Projected Owner Group | 13 |
| ASPECT | R9. | Aspect of Plot Center | 13 |
| POSITION | R10. | Position on Slope | 13 |
| SLOPE | R11. | Slope Angle at Plot Center | 13 |
| PHYCLS | R12. | Physiographic Class | 13 |
| STDORG | R13. | Stand Origin | 13 |
| SEEDSOR | R14. | Seed source | 13 |
| CUNCOND | R15. | Conifer Understory Condition | 13 |
| CUNSPP | R16. | Conifer Understory Species | 13 |
| BACUR | R17. | Basal Area Current Value | 13 |
| BAORG | R18. | Basal Area Original Value | 13 |
| BAPRJ | R19. | Basal Area Projected Value | 14 |
| FORTCUR | C4. | Current Forest Type | 14 |
| FORTPRJ | R20. | Projected Forest Type | 14 |
| SZDNCUR | R21. | Current Stand Size-Density | 14 |


[1/] C = Common Information Record (CIR) variable
R = Record variable.

RECORD 2 (PLOTFOR) -(continued)

| Variable name | Variable type & number[1/] | Definition | Page |
|---|---|---|---|
| SZDNPRJ | R22. | Projected Stand Size-Density | 14 |
| FORTORG | R23. | Original Forest Type | 14 |
| SZDNORG | R24. | Original Stand Size-Density | 14 |
| STDAGE | R25. | Stand Age | 14 |
| PRJAGE | R26. | Projected Stand Age | 14 |
| SITECLS | R27. | Site Class | 14 |
| SITEIND | C5. | Site Index | 14 |
| SISPP | R28. | Site Index Species | 14 |
| EXPVOL | R29. | Volume Expander | 14 |
| GSCUVS | R30. | Growing Stock CuFt Volume Softwoods | 15 |
| GSCUVH | R31. | Growing Stock CuFt Volume Hardwoods | 15 |
| PRJSVOL | R32. | Projected G.S. CuFt Volume Softwoods | 15 |
| PRJHVOL | R33. | Projected G.S. CuFt Volume Hardwoods | 15 |
| STDAREA | R34. | Stand Area | 15 |
| AREACOND | R35. | Stocking Level Class | 15 |
| GRSTKPC | R36. | Growing Stock Tree Stocking | 15 |
| ROUGHPC | R37. | Rough Tree Stocking | 16 |
| ROTENPC | R38. | Rotten Tree Stocking | 16 |
| PRJGSPC | R39. | Projected G.S. Tree Stocking | 16 |
| PRJRGPC | R40. | Projected Rough Tree Stocking | 16 |
| PRJRTPC | R41. | Projected Rotten Tree Stocking | 16 |
| SG01 | C8. | Presence or Absence of Jack Pine | 15 |
| SG02 | C9. | Presence or Absence of Red Pine | 15 |
| SG03 | C10. | Presence or Absence of White Pine | 15 |
| SG04 | C11. | Presence or Absence of Ponderosa Pine | 15 |
| SG05 | C12. | Presence or Absence of Loblolly Pine | 15 |
| SG06 | C13. | Presence or Absence of Shortleaf Pine | 15 |
| SG07 | C14. | Presence or Absence of Other Yellow Pines | 15 |
| SG08 | C15. | Presence or Absence of White Spruce | 15 |
| SG09 | C16. | Presence or Absence of Black Spruce | 15 |
| SG10 | C17. | Presence or Absence of Balsam Fir | 15 |
| SG11 | C18. | Presence or Absence of Hemlock | 15 |
| SG12 | C19. | Presence or Absence of Tamarack | 15 |
| SG13 | C20. | Presence or Absence of Bald Cypress | 15 |
| SG14 | C21. | Presence or Absence of E. Red Cedar | 15 |
| SG15 | C22. | Presence or Absence of N. white-cedar | 15 |
| SG16 | C23. | Presence or Absence of (not assigned) | 15 |
| SG17 | C24. | Presence or Absence of Other Softwoods | 15 |
| SG18 | C25. | Presence or Absence of Select White Oak | 15 |
| SG19 | C26. | Presence or Absence of Other White Oak | 15 |
| SG20 | C27. | Presence or Absence of Select Red Oak | 15 |
| SG21 | C28. | Presence or Absence of Other Red Oak | 16 |
| SG22 | C29. | Presence or Absence of Select Hickory | 16 |
| SG23 | C30. | Presence or Absence of Other Hickory | 16 |
| SG24 | C31. | Presence or Absence of Basswood | 16 |
| SG25 | C32. | Presence or Absence of Beech | 16 |
| SG26 | C33. | Presence or Absence of Yellow Birch | 16 |
| SG27 | C34. | Presence or Absence of Hard Maple | 16 |
| SG28 | C35. | Presence or Absence of Soft Maple | 16 |
| SG29 | C36. | Presence or Absence of Elm | 16 |
| SG30 | C37. | Presence or Absence of Black Ash | 16 |
| SG31 | C38. | Presence or Absence of White-Green Ash | 16 |
| SG32 | C39. | Presence or Absence of Sycamore | 16 |
| SG33 | C40. | Presence or Absence of Cottonwood | 16 |
| SG34 | C41. | Presence or Absence of Willow | 16 |
| SG35 | C42. | Presence or Absence of Hackberry | 16 |
| SG36 | C43. | Presence or Absence of Balsam Poplar | 16 |
| SG37 | C44. | Presence or Absence of Bigtooth Aspen | 16 |
| SG38 | C45. | Presence or Absence of Quaking Aspen | 16 |
| SG39 | C46. | Presence or Absence of Paper Birch | 16 |
| SG40 | C47. | Presence or Absence of River Birch | 16 |
| SG41 | C48. | Presence or Absence of Sweetgum | 16 |
| SG42 | C49. | Presence or Absence of Blackgum and Tupelo | 16 |
| SG43 | C50. | Presence or Absence of Black Cherry | 16 |

[1/] C = Common Information Record (CIR) variable
R = Record variable

RECORD 2 (PLOTFOR) -(continued)

| Variable name | Variable type & number[1/] | Definition | Page |
|---|---|---|---|
| SG44 | C51. | Presence or Absence of Black Walnut | 16 |
| SG45 | C52. | Presence or Absence of Butternut | 16 |
| SG46 | C53. | Presence or Absence of Yellow Poplar | 16 |
| SG47 | C54. | Presence or Absence of (not assigned) | 16 |
| SG48 | C55. | Presence or Absence of (not assigned) | 16 |
| SG49 | C56. | Presence or Absence of Other Hardwoods | 16 |
| SG50 | C57. | Presence or Absence of Noncommercial | 16 |

RECORD 3 (TREE)

| Variable name | Variable type & number[1/] | Definition | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 17 |
| TRNUM | R1. | Tree Number | 17 |
| TRHIST | R2. | Tree History-Status | 17 |
| PRJHIST | R3. | Projected Tree History-Status | 17 |
| SPECIES | R4. | Tree Species | 17 |
| DBHORG | R5. | Original Diameter Breast High | 17 |
| DBHCUR | R6. | Current Diameter Breast High | 17 |
| DBHPRJ | R7. | Projected Diameter Breast High | 17 |
| SOUND | R8. | Soundness Code | 17 |
| LOGGR | R9. | Log Grade | 18 |
| CAVIT | R10. | Cavities Code | 18 |
| CRRAT | R11. | Crown Ratio | 18 |
| PRJCRRAT | R12. | Projected Crown Ratio | 18 |
| CRCLS | R13. | Crown Class | 18 |
| DAMDETH | R14. | Damage Cause of Death | 18 |
| TRCLS | R15. | Tree Class | 18 |
| PRJTRCLS | R16. | Projected Tree Class | 18 |
| PTOCC | R17. | Point Occupancy | 18 |
| EXPANDR | R18. | Number of Trees Per Acre | 18 |
| PRJEXPD | R19. | Projection Factor | 18 |
| MORTFAC | R20. | Mortality Factor | 18 |
| NCSPGP | R21. | North Central Species Group | 19 |
| TRSZCLS | R22. | Tree Size Class | 19 |
| PRJTRSZ | R23. | Projected Tree Size Class | 19 |
| TRSIND | R24. | Tree Site Index | 19 |
| NETCUVL | R25. | Net Cubic Foot Volume in Tree | 19 |
| PRJNCVL | R26. | Projected Net CuFt Volume in Tree | 19 |

RECORD 4 (SHRUBT)

| | | | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 20 |
| SBTNUM | R1. | Shrub Number-Tall | 20 |
| SHRUBSP | R2. | Shrub Species-Tall | 20 |
| DIACLS | R3. | Diameter Class | 20 |
| COUNT | R4. | Count of Number of Individuals | 20 |
| EXPSHUT | R5. | Expander for Shrub Tall | 20 |

RECORD 5 (SHRUBL)

| | | | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 20 |
| SBLNUM | R1. | Shrub Number-Low | 20 |
| SHUSPL | R2. | Shrub Species-Low | 20 |
| COVER | R3. | Percent Ground Cover | 21 |
| EXPSHUL | R4. | Expander for Low Shrub | 21 |

RECORD 6 (SITETR)

| | | | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 21 |
| TRNUM | R1. | Tree Number | 21 |
| DBHSI | R2. | Diameter Breast High Site Index Tre | 21 |
| SPPSI | R3. | Species Site Index Tree | 21 |
| HTSI | R4. | Height Site Index Tree | 21 |
| AGESI | R5. | Total Age Site Index Tree | 21 |
| YRSADD | R6. | Years Added to DBH Age for Total Ag | 21 |

[1/] C = Common Information Record (CIR) variable
R = Record variable.

RECORD 7 (POINT)

| Variable name | Variable type & number[1/] | Definition | Page |
|---|---|---|---|
| IDENT | C6. | Case Identifier | 22 |
| PTNUM | R1. | Point Number | 22 |
| PTCLS | R2. | Point Class | 22 |
| INDEX | | | 23 |

1/C = Common Information Record (CIR) variable
R = Record variable.

This data base contains data for an entire State or a portion of State. The code book is the same for the entire State. Each case contains the data from one survey plot in seven record types. Record types 1 and 2 contain plot level data and record types 3 through 7 contain tree and shrub data.

*** RECORD 1 (PLOTALL) DEFINITION ***

Record type 1 (PLOTALL) contains data gathered for all plots.

```
MAXIMUM RECS/CASE:   1
CASE IDENTIFIER:     IDENT (A)

C6.     IDENT    , Case Identifier
                   (XYYYYY where;
                    X = FIA unit number
                    YYYYY = Plot number.
                    Every plot within a data base
                    has a unique case identifier.
                    See appendix for unit codes.)
                FORMAT:  I6
             DATA TYPE:  I*4

R1.     STATE    , State Code
                FORMAT:  I2
             DATA TYPE:  I*2
           VALUE LABELS: (17) Illinois   (26) Michigan    (38) North Dakota
                         (18) Indiana    (27) Minnesota   (46) South Dakota
                         (19) Iowa       (29) Missouri    (55) Wisconsin
                         (20) Kansas     (31) Nebraska

C1.     UNTCO    , Unit and County
                   (XYY where;
                    X = Unit Number
                    YY = County number.
                    See appendix for codes)
                FORMAT:  I3
             DATA TYPE:  I*2

R2.     SAMPKND  , Sample Kind
                FORMAT:  I1
             DATA TYPE:  I*1
           VALUE LABELS:
              (1) New 10-Point Full              (6) Remeasurement Partial
              (2) Remeasurement                  (7) New 10-Point Partial
              (3) Remeasurement NF               (8) Remeasurement Partial
              (4) New 10-Point Full NF           (9) Remeasurement Updated
              (5) Reserve[Phantom]               (0) Stereo dots [not gr.ck]

R3.     DISTURB  , Disturbance Code
                    Not used prior to 1982
                FORMAT:  I1
             DATA TYPE:  I*1
           VALUE LABELS:
              (1) Forest to forest undisturbed   (4) Forest to nonforest
              (2) Nonforest to forest disturbed  (5) Nonforest to nonforest
              (3) Nonforest to forest

R4.     PICODE   , Photo Interpretation Code
                   (See appendix for codes)
                FORMAT:  I2
             DATA TYPE:  I*2
```

R5. PISTRAT , Photo Stratification Code
(See appendix for codes)
FORMAT: I2
DATA TYPE: I*2

C2. GLUCUR , Ground Land Use Current
R6. GLUORG , Ground Land Use Original
R7. GLUPRJ , Ground Land Use Projected
FORMAT: I2
DATA TYPE: I*2
VALUE LABELS:

| | |
|---|---|
| (20) Commercial Forest | (21) Pastured Com. Fore: |
| (22) Plantation | (40) Unproductive Forest |
| (41) Reserved Forest | (45) Reserved For.Prod. |
| (46) Christmas Tr.Plant. | (51) Cropland With Trees |
| (52) Pasture With Trees | (53) Wooded Strip |
| (54) Idle Farmland W-T | (55) Marsh With Trees |
| (56) Narrow Windbreaks | (57) Wide Windbreaks |
| (58) Windbreaks | (59) Wooded Pasture |
| (61) Cropland Wo-T | (62) Improved Pasture Wo- |
| (64) Idle Farmland Wo-T | (65) Marsh Wo-T |
| (66) Other Farmland | (67) Urban and Other |
| (68) Rights-of-way | (69) Nonforest Reserve |
| (71) Urban For. Reserved | (72) Urban With Trees |
| (80) Noncensus Water | (90) Census Water |

R8. CHANGE , Reason for GLU change
FORMAT: I1
DATA TYPE: I*1

R9. UTREND , Use Trend
See appendix for deff.
FORMAT: I3
DATA TYPE: I*2

R10. STDHIST , Stand History
See appendix for deff.
FORMAT: I2
DATA TYPE: I*2

R11. PRJSTDH , Projected Stand History
See appendix for deff.
FORMAT: I2
DATA TYPE: I*2

R12. PHOTAGE , Photo Age
(years)
FORMAT: I1
DATA TYPE: I*1

R13. SURDATE , Survey Date
FORMAT: A4 WITH DATE MAP: 'MMYY'
DATA TYPE: I*4

R14. PRJDATE , Projected Date
FORMAT: A4 WITH DATE MAP: 'MMYY'
DATA TYPE: I*4

R15. GROPER , Remeasurement Peroid
(years)
FORMAT: I2
DATA TYPE: I*2

R16. USESET , Use Setting-Recreation
Not used prior to 1982
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS:

| | |
|---|---|
| (1) Primitive | (2) Semi-prim. non-mot |
| (3) Semi-prim. motor | (4) Roaded natural |
| (5) Rural | (6) Urban |

R17. USEAREA , Use Area-Recreation
Not used prior to 1982
FORMAT: I1
DATA TYPE: I*1

R18. USEPOST , Use Posted-Recreation
Not used prior to 1982
FORMAT: I1
DATA TYPE: I*1

R19. SOILCLS , Soil Classification
Available for future use
FORMAT: I8
DATA TYPE: I*4

C7. EXPAREA , Area Expander
(acres)
FORMAT: I5
DATA TYPE: I*4

R20. PRJAREA1, Projected Area Expander 1
(acres)
FORMAT: I5
DATA TYPE: I*4

R21. PRJAREA2, Projected Area Expander 2
(acres)
FORMAT: I5
DATA TYPE: I*4

R22. UTMZONE , UTM Zone Code
FORMAT: I2
DATA TYPE: I*2

R23. UTMNORTH, UTM Northing To Nearest 1000 Meters
FORMAT: I5
DATA TYPE: I*4

R24. UTMEAST , UTM Easting To Nearest 1000 Meters
FORMAT: I5
DATA TYPE: I*4

R25. TOWNSHIP, Township
FORMAT: A4
DATA TYPE: A*6

R26. RANGE , Range
FORMAT: A4
DATA TYPE: A*6

R27. SECTION , Section
FORMAT: I2
DATA TYPE: I*2

R28. EXPREMEA, Remeasurement Expander
(acres)
FORMAT: I6
DATA TYPE: I*4

RECORD 2 (PLOTFOR ) DEFINITION ***

Record type 2 (PLOTFOR) contains data for plots classified as forest, at either measurement occasion for remeasurement plots, or for plots classified as non-forest with trees.

```
MAXIMUM RECS/CASE:    1
CASE IDENTIFIER:      IDENT (A)

C6.    IDENT   , Case Identifier
                 (Unit & Seq.Plot Num.)
             FORMAT:  I6
          DATA TYPE:  I*4

R1.    DWT     , Distance to Water Type
                 Not used prior to 1974
             FORMAT:  I1
          DATA TYPE:  I*1
       VALUE LABELS:  (1) Streams          (2) Lakes
                      (3) Swamps           (4) Farm ponds

R2.    DWS     , Distance to Water Size of Water
                 (acres)
                 Not used prior to 1982
             FORMAT:  I3
          DATA TYPE:  I*2

R3.    DWV     , Distance to Water Value
                 (chains)
                 Not used prior to 1974
             FORMAT:  I4
          DATA TYPE:  I*2

R4.    DRT     , Distance to Road Type of Road
                 Not used prior to 1974
             FORMAT:  I1
          DATA TYPE:  I*1
       VALUE LABELS:  (1) Paved-4 lane     (2) Paved-2 lane
                      (3) Improved-gravel

R5.    DRV     , Distance to Road Value
                 (chains)
                 Not used prior to 1974
             FORMAT:  I4
          DATA TYPE:  I*2

R6.    NATFOR  , National Forest Code
             FORMAT:  I3
          DATA TYPE:  I*2
       VALUE LABELS:  (902) Chequamegon    (903) Chippewa
                      (904) Huron-Manistee (906) Nicolet
                      (907) Ottawa         (908) Shawnee
                      (909) Superior       (910) Hiawatha
                      (911) Wayne-Hoosier  (918) Mark Twain

R7.    RANDIST , Ranger District
             FORMAT:  I2
          DATA TYPE:  I*2

C3.    OWNGP   , Ownership Group
             FORMAT:  I1
          DATA TYPE:  I*1
       VALUE LABELS:  (1) National forest  (2) Other public
                      (3) Private
```

R8. PRJOWN , Projected Owner Group
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) National forest (2) Other public
(3) Private

R9. ASPECT , Aspect of Plot Center
(degrees)
Not used prior to 1973
FORMAT: I3
DATA TYPE: I*2

R10. POSITION, Position on Slope
Not used prior to 1973
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) Top quarter (2) Upper quarter
(3) Lower quarter (4) Level or lowest quarter

R11. SLOPE , Slope Angle at Plot Center
(percent)
Not used prior to 1973
FORMAT: I2
DATA TYPE: I*2

R12. PHYCLS , Physiographic Class
(Degree of Wet-Dryness)
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (3) Xeric (4) Xeromesic
(5) Mesic (6) Hydromesic
(7) Hydric

R13. STDORG , Stand Origin
(Planted-Natural)

FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) Natural (2) GE 40 percent artificial
(3) LT 40 percent artificial

R14. SEEDSOR , Seed Source
(Degree of Conifer Seed Source)
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS (1) Adequate softwood (2) Adequate hardwood
(3) Adeq.soft & hard (4) Inadequate all spp.

R15. CUNCOND , Conifer Understory Condition
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) None or inadequate (2) Planted adequate
(3) Planted needs treat. (4) Natural adequate
(5) Natural needs treat.

R16. CUNSPP , Conifer Understory Species
(See appendix for codes)
FORMAT: I3
DATA TYPE: I*2

R17. BACUR , Basal Area Current Value
(sq feet)
FORMAT: I3
DATA TYPE: I*2

R18. BAORG , Basal Area Original Value
(sq feet)
FORMAT: I3
DATA TYPE: I*2

R19. BAPRJ , Basal Area Projected Value
(sq feet)
FORMAT: I3
DATA TYPE: I*2

C4. FORTCUR , Current Forest Type
(See appendix for codes)
FORMAT: I2
DATA TYPE: I*2

R20. FORTPRJ , Projected Forest Type
(See appendix for codes)
FORMAT: I2
DATA TYPE: I*2

R21. SZDNCUR , Current Stand Size-Density

R22. SZDNPRJ , Projected Stand Size-Density
FORMAT: I2
DATA TYPE: I*2
VALUE LABELS: (11) Sawtimber-Good (21) Poletimber-Good
(12) Sawtimber-Medium (22) Poletimber-Medium
(13) Sawtimber-Poor (23) Poletimber-Poor
(31) Seed-Saps-Good (40) Nonstocked
(32) Seed-Saps-Medium
(33) Seed-Saps-Poor

R23. FORTORG , Original Forest Type
(See appendix for codes)
FORMAT: I2
DATA TYPE: I*2

R24. SZDNORG , Original Stand Size-Density
FORMAT: I2
DATA TYPE: I*2
VALUE LABELS: (10) Sawtimber (20) Poletimber
(30) Sapling & seedling (40) Nonstocked

R25. STDAGE , Stand Age
(years)
FORMAT: I3
DATA TYPE: I*2

R26. PRJAGE , Projected Stand Age
(years)
FORMAT: I3
DATA TYPE: I*2

R27. SITECLS , Site Class
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) 225+ cu.ft.
(2) 165 to 225 cu.ft.
(3) 120 to 165 cu.ft.
(4) 85 to 120 cu.ft.
(5) 50 to 85 cu.ft.
(6) Less than 50 cu.ft.

C5. SITEIND , Site Index
(feet at age 50)
FORMAT: I2
DATA TYPE: I*2

R28. SISPP , Site Index Species
(See appendix for codes)
FORMAT: I3
DATA TYPE: I*2

R29. EXPVOL , Volume Expander
(acres)
FORMAT: I5
DATA TYPE: I*4

R30. GSCUVS , Growing Stock CuFt Volume Softwoods
(cubic feet per acre)
FORMAT: I6
DATA TYPE: I*4

R31. GSCUVH , Growing Stock CuFt Volume Hardwoods
(cubic feet per acre)
FORMAT: I6
DATA TYPE: I*4

R32. PRJSVOL , Projected G.S. CuFt Volume Softwoods
(cubic feet per acre)
FORMAT: I6
DATA TYPE: I*4

R33. PRJHVOL , Projected G.S. CuFt Volume Hardwoods
(cubic feet per acre)
FORMAT: I6
DATA TYPE: I*4

R34. STDAREA , Stand Area
(acres)
FORMAT: I3
DATA TYPE: I*2

R35. AREACOND, Stocking Level Class
(See appendix for codes)
FORMAT: I2
DATA TYPE: I*2

R36. GRSTKPC , Growing Stock Tree Stocking
(percent)
FORMAT: I3
DATA TYPE: I*2

Variables C8 (SG01) through C57 (SG50) indicate the presence or absence of a particular species group on the plot. For example, SG01 indicates the presence or absence of jack pine on the plot. If SG01 is 0, then no jack pine was found on the plot. SG01 is 1 when jack pine seedling and/or sapling trees were found, but jack pine poletimber or sawtimber was not. SG01 is 2, jack pine poletimber, but not sawtimber, was present; if SG01 is 3, jack pine sawtimber was present.

C8. SG01 , Presence or Absence of Jack Pine
C9. SG02 , Presence or Absence of Red Pine
C10. SG03 , Presence or Absence of White Pine
C11. SG04 , Presence or Absence of Ponderosa Pine
C12. SG05 , Presence or Absence of Loblolly Pine
C13. SG06 , Presence or Absence of Shortleaf Pine
C14. SG07 , Presence or Absence of Other Yellow Pine
C15. SG08 , Presence or Absence of White Spruce
C16. SG09 , Presence or Absence of Black Spruce
C17. SG10 , Presence or Absence of Balsam Fir
C18. SG11 , Presence or Absence of Hemlock
C19. SG12 , Presence or Absence of Tamarack
C20. SG13 , Presence or Absence of Bald Cypress
C21. SG14 , Presence or Absence of E. Red Cedar
C22. SG15 , Presence or Absence of N. White-cedar
C23. SG16 , Presence or Absence of (not assigned)
C24. SG17 , Presence or Absence of Other Softwoods
C25. SG18 , Presence or Absence of Select White Oak
C26. SG19 , Presence or Absence of Other White Oak
C27. SG20 , Presence or Absence of Select Red Oak

C28. SG21 , Presence or Absence of Other Red Oak
C29. SG22 , Presence or Absence of Select Hickory
C30. SG23 , Presence or Absence of Other Hickory
C31. SG24 , Presence or Absence of Basswood
C32. SG25 , Presence or Absence of Beech
C33. SG26 , Presence or Absence of Yellow Birch
C34. SG27 , Presence or Absence of Hard Maple
C35. SG28 , Presence or Absence of Soft Maple
C36. SG29 , Presence or Absence of Elm
C37. SG30 , Presence or Absence of Black Ash
C38. SG31 , Presence or Absence of White-Green Ash
C39. SG32 , Presence or Absence of Sycamore
C40. SG33 , Presence or Absence of Cottonwood
C41. SG34 , Presence or Absence of Willow
C42. SG35 , Presence or Absence of Hackberry
C43. SG36 , Presence or Absence of Balsam Poplar
C44. SG37 , Presence or Absence of Bigtooth Aspen
C45. SG38 , Presence or Absence of Quaking Aspen
C46. SG39 , Presence or Absence of Paper Birch
C47. SG40 , Presence or Absence of River Birch
C48. SG41 , Presence or Absence of Sweet Gum
C49. SG42 , Presence or Absence of Black Gum and Tupelo
C50. SG43 , Presence or Absence of Black Cherry
C51. SG44 , Presence or Absence of Black Walnut
C52. SG45 , Presence or Absence of Butternut
C53. SG46 , Presence or Absence of Yellow Poplar
C54. SG47 , Presence or Absence of (not assigned)
C55. SG48 , Presence or Absence of (not assigned)
C56. SG49 , Presence or Absence of Other Hardwoods
C57. SG50 , Presence or Absence of Noncommercial
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) Sapling & seedling (2) Poletimber (3) Sawtimber

R37. ROUGHPC , Rough Tree Stocking
(percent)
FORMAT: I3
DATA TYPE: I*2

R38. ROTENPC , Rotten Tree Stocking
(percent)
FORMAT: I3
DATA TYPE: I*2

R39. PRJGSPC , Projected G.S. Tree Stocking
(percent)
FORMAT: I3
DATA TYPE: I*2

R40. PRJRGPC , Projected Rough Tree Stocking
(percent)
FORMAT: I3
DATA TYPE: I*2

R41. PRJRTPC , Projected Rotten Tree Stocking
(percent)
FORMAT: I3
DATA TYPE: I*2

*** RECORD 3 (TREE) DEFINITION ***

Record type 3 (TREE) contains the individual tree data tallied on each plot. The species group variable is set at 1, 2, or 3 on the CIR if a species group is present on the plot; otherwise it is blank or missing, indicating that the species is not present.

```
MAXIMUM RECS/CASE:   250
CASE IDENTIFIER:     IDENT (A)
SORT IDENTIFIER: (1) TRNUM (A)

C6.    IDENT    , Case Identifier
                  (Unit & Seq.Plot Num.)
              FORMAT:  I6
           DATA TYPE:  I*4

R1.    TRNUM    , Tree Number
              FORMAT:  I4
           DATA TYPE:  I*2

R2.    TRHIST   , Tree History-Status
                  On remeasurement plots the first
                  digit is original status, the
                  second digit is current status
              FORMAT:  I2
           DATA TYPE:  I*2
         VALUE LABELS: (0) No status            (1) Growing stock live
                       (2) Cull live            (3) Ingrowth
                       (4) Dead[salvable]       (5) Dead[mortality]
                       (6) Ongrowth             (7) Stump[salvaged]
                       (8) Stump[utilized]      (9) Stump[not utilized]

R3.    PRJHIST , Projected Tree History-Status
              FORMAT:  I1
           DATA TYPE:  I*1
         VALUE LABELS: (0) No status            (1) Growing stock live
                       (2) Cull live            (3) Ingrowth
                       (5) Dead[mortality]      (8) Stump[utilized]

R4.    SPECIES , Tree Species
                  (See appendix for codes)
              FORMAT:  I3
           DATA TYPE:  I*2

R5.    DBHORG   , Original Diameter Breast High
                  (inches)
              FORMAT:  D4.2
           DATA TYPE:  R*8

R6.    DBHCUR   , Current Diameter Breast High
                  (inches)
              FORMAT:  D4.2
           DATA TYPE:  R*8

R7.    DBHPRJ   , Projected Diameter Breast High
                  (inches)
              FORMAT:  D4.2
           DATA TYPE:  R*8

R8.    SOUND    , Soundness Code
              FORMAT:  I1
           DATA TYPE:  I*1

R9.    LOGGR    , Log Grade
              FORMAT:  I1
           DATA TYPE:  I*1
```

R10. CAVIT , Cavities Code not used prior to 1981
(See appendix for codes)
FORMAT: I2
DATA TYPE: I*2

R11. CRRAT , Crown Ratio
(percent live crown)
FORMAT: I1
DATA TYPE: I*1

R12. PRJCRRAT, Projected Crown Ratio
(percent live crown)
FORMAT: I1
DATA TYPE: I*1

R13. CRCLS , Crown Class
FORMAT: I1
DATA TYPE: I*1
VALUE LABELS: (1) Open grown (2) Dominant
(3) Codominant (4) Intermediate
(5) Overtopped

R14. DAMDETH , Damage Cause of Death
FORMAT: I2
DATA TYPE: I*2
VALUE LABELS: (0) None (10) Insect
(11) Poplar borer (20) Disease
(21) Phellinus tremulae (22) Hypoxylon canker
(23) Other cankers (24) Blister rust
(25) Dwarf mistletoe (26) Shoot blights
(27) Butternut canker (28) Nectria canker
(29) Eutypella canker (30) Fire
(40) Animal (41) Beaver
(42) Cattle (43) Deer
(44) Porcupine (45) Sapsuckers
(50) Weather (60) Suppression
(70) Unknown (71) Missing or dead top
(72) Poor form (80) Logging
(81) Logging (82) Timber stand imp.
(84) Land clearing (85) Conversion

R15. TRCLS , Tree Class
R16. PRJTRCLS, Projected Tree Class
FORMAT: I2
DATA TYPE: I*2
VALUE LABELS: (10) Growing stock des. (20) Growing stock acc.
(30) Cull rough (31) Cull short-log
(40) Cull rotten

R17. PTOCC , Point Occupancy
FORMAT: I1
DATA TYPE: I*1

R18. EXPANDR , Number of Trees Per Acre
FORMAT: D4.2
DATA TYPE: R*8

R19. PRJEXPD , Projection Factor
(num. trees per acre)
FORMAT: D4.2
DATA TYPE: R*8

R20. MORTFAC , Mortality Factor
(num. trees per acre per year)
FORMAT: D4.2
DATA TYPE: R*8

R21. NCSPGP , North Central Species Group

FORMAT: I2

VALUE LABELS:

| | |
|---|---|
| (1) Jack pine | (2) Red pine |
| (3) White pine | (4) Ponderosa pine |
| (5) Loblolly pine | (6) Shortleaf pine |
| (7) Other yellow pines | (8) White spruce |
| (9) Black spruce | (10) Balsam fir |
| (11) Hemlock | (12) Tamarack |
| (13) Bald cypress | (14) Eastern redcedar |
| (15) Northern white-cedar | (16) |
| (17) Other softwoods | (18) Select white oak |
| (19) Other white oak | (20) Select red oak |
| (21) Other red oak | (22) Select hickory |
| (23) Other hickory | (24) Basswood |
| (25) Beech | (26) Yellow birch |
| (27) Hard maple | (28) Soft maple |
| (29) Elm | (30) Black ash |
| (31) White & green ash | (32) Sycamore |
| (33) Cottonwood | (34) Willow |
| (35) Hackberry | (36) Balsam poplar |
| (37) Bigtooth aspen | (38) Quaking aspen |
| (39) Paper birch | (40) River birch |
| (41) Sweetgum | (42) Blackgum and tupelo |
| (43) Black cherry | (44) Black walnut |
| (45) Butternut | (46) Yellow poplar |
| (47) | (48) |
| (49) Other hardwoods | (50) Noncommercial spp. |

R22. TRSZCLS , Tree Size Class

FORMAT: I1

DATA TYPE: I*1

VALUE LABELS: (1) Sapling & seedling (2) Poletimber
(3) Sawtimber

R23. PRJTRSZ , Projected Tree Size Class

FORMAT: I1

DATA TYPE: I*1

VALUE LABELS: (1) Seeds & saps
(2) Poletimber
(3) Sawtimber

R24. TRSIND , Tree Site Index
(feet age 50)

FORMAT: I2

DATA TYPE: I*2

R25. NETCUVL , Net Cubic Foot Volume in Tree
(cubic feet per acre)

FORMAT: I6

DATA TYPE: I*4

R26. PRJNCVL , Projected Net CuFt Volume in Tree
(cubic feet per acre)

FORMAT: I6

DATA TYPE: I*4

RECORD 4 (SHRUBT) DEFINITION ***

Record type 4 (SHURBT) contains the data tallied for tall woody shrubs on forest plots.

```
MAXIMUM RECS/CASE:   100
CASE IDENTIFIER:     IDENT (A)
SORT IDENTIFIER: (1) SBTNUM (A)

C6.     IDENT    , Case Identifier
                   (Unit & Seq.Plot Num.)
                FORMAT:  I6
             DATA TYPE:  I*4

R1.     SBTNUM   , Shrub Number-Tall
                FORMAT:  I4
             DATA TYPE:  I*2

R2.     SHRUBSP  , Shrub Species-Tall
                   (See appendix for codes)
                FORMAT:  I3
             DATA TYPE:  I*2

R3.     DIACLS   , Diameter Class
                   (inches)
                FORMAT:  I2
             DATA TYPE:  I*2
           VALUE LABELS: (1) 0.0-0.19        (2) 0.2-0.29
                         (3) 0.3-0.39        (4) 0.4-0.49
                         (5) 0.5-0.99        (10) 1.0-1.49
                         (15) 1.5-1.99       (20) 2.0-2.49
                         (25) 2.5-2.99       (30) 3.0-3.49
                              etc.

R4.     COUNT    , Count of Number of Individuals
                FORMAT:  I3
             DATA TYPE:  I*2

R5.     EXPSHUT  , Expander for Shrub Tall
                   (shrubs per acre)

                FORMAT:  D4.2
             DATA TYPE:  R*8
```

*** RECORD 5 (SHRUBL) DEFINITION ***

Record type 5 (SHRUBL) contains the data tallied for low herbaceous shrubs on forest plots.

```
MAXIMUM RECS/CASE:   30
CASE IDENTIFIER:     IDENT (A)

SORT IDENTIFIER: (1) SBLNUM (A)

C6.     IDENT    , Case Identifier
                   (Unit & Seq.Plot Num.)
                FORMAT:  I6
             DATA TYPE:  I*4

R1.     SBLNUM   , Shrub Number-Low
                FORMAT:  I4
             DATA TYPE:  I*2

R2.     SHUSPL   , Shrub Species-Low
                   (See appendix for codes)
                FORMAT:  I3
             DATA TYPE:  I*2
```

```
R3.     COVER    , Percent Ground Cover
                FORMAT:  I1
            DATA TYPE:  I*1
          VALUE LABELS:  (1) Solitary Plant        (2) Few Plants LT 5 percent
                         (3) Numerous Plants       (4) 5-25 percent
                         (5) 25-50 percent         (6) 50-90 percent
                         (7) 90-100 percent

R4.     EXPSHUL , Expander for Low Shrub
                  (shrubs per acre)
                FORMAT:  D4.2
            DATA TYPE:  R*8
```

*** RECORD 6 (SITETR) DEFINITION ***

Record type 6 (SITETR) contains data tallied for site trees on forest plots.

```
MAXIMUM RECS/CASE:   10
CASE IDENTIFIER:     IDENT (A)

SORT IDENTIFIER: (1) TRNUM (A)

C6.     IDENT    , Case Identifier
                  (Unit & Seq.Plot Num.)
                FORMAT:  I6
            DATA TYPE:  I*4

R1.     TRNUM    , Tree Number
                FORMAT:  I4
            DATA TYPE:  I*2

R2.     DBHSI    , Diameter Breast High Site Index Tree
                  (inches)
                FORMAT:  D3.1
            DATA TYPE:  R*8

R3.     SPPSI    , Species Site Index Tree
                  (See appendix for codes)
                FORMAT:  I3
            DATA TYPE:  I*2

R4.     HTSI     , Height Site Index Tree
                  (feet)
                FORMAT:  I3
            DATA TYPE:  I*2

R5.     AGESI    , Total Age Site Index Tree
                  (years)
                FORMAT:  I3
            DATA TYPE:  I*2

R6.     YRSADD   , Years Added to DBH Age for Total Age
                FORMAT:  I2
            DATA TYPE:  I*2
```

RECORD 7 (POINT) DEFINITION ***

Record type 7 (POINT) contains data tallied for nonstocked points on forest plots.

```
MAXIMUM RECS/CASE:   10
CASE IDENTIFIER:     IDENT (A)

SORT IDENTIFIER: (1) PTNUM (A)

C6.     IDENT    , Case Identifier
                   (Unit & Seq.Plot Num.)
                FORMAT:  I6
             DATA TYPE:  I*4

R1.     PTNUM    , Point Number
                FORMAT:  I4
             DATA TYPE:  I*2

R2.     PTCLS    , Point Class
                FORMAT:  I2
             DATA TYPE:  I*2
           VALUE LABELS: (50) Inhibiting Vegetation
                         (51) Inhib. Veg.-Grass
                         (52) Inhib. Veg.-Shrubs
                         (53) Inhib. Veg.-Vines
                         (54) Inhib. Veg.-Other
                         (60) Nonstocked
                         (70) Overtopped
                         (80) Nonstockable
                         (81) Nonstockable-Rocks
                         (82) Nonstockable-Water
                         (83) Nonstockable-Other
```

INDEX


| Variable name | Record number | Variable number | Definition | Page |
|---|---|---|---|---|
| AGESI | 6 | R5. | Total Age Site Index Tree | 21 |
| AREACOND | 2 | R35. | Stocking Level Class | 15 |
| ASPECT | 2 | R9. | Aspect of Plot Center | 13 |
| BACUR | 2 | R17. | Basal Area Current Value | 13 |
| BAORG | 2 | R18. | Basal Area Original Value | 13 |
| BAPRJ | 2 | R19. | Basal Area Projected Value | 14 |
| CAVIT | 3 | R10. | Cavities Code | 18 |
| CHANGE | 1 | R8. | Reason for GLU change | 10 |
| COUNT | 4 | R4. | Count of Number of Individuals | 20 |
| COVER | 5 | R3. | Percent Ground Cover | 21 |
| CRCLS | 3 | R13. | Crown Class | 18 |
| CRRAT | 3 | R11. | Crown Ratio | 18 |
| CUNCOND | 2 | R15. | Conifer Understory Condition | 13 |
| CUNSPP | 2 | R16. | Conifer Understory Species | 13 |
| DAMDETH | 3 | R14. | Damage Cause of Death | 18 |
| DBHCUR | 3 | R6. | Current Diameter Breast High | 17 |
| DBHORG | 3 | R5. | Original Diameter Breast High | 17 |
| DBHPRJ | 3 | R7. | Projected Diameter Breast High | 17 |
| DBHSI | 6 | R2. | Diameter Breast High Site Index Tree | 21 |
| DIACLS | 4 | R3. | Diameter Class | 20 |
| DISTURB | 1 | R3. | Disturbance Code | 9 |
| DRT | 2 | R4. | Distance to Road Type of Road | 12 |
| DRV | 2 | R5. | Distance to Road Value | 12 |
| DWS | 2 | R2. | Distance to Water Size of Water | 12 |
| DWT | 2 | R1. | Distance to Water Type | 12 |
| DWV | 2 | R3. | Distance to Water Value | 12 |
| EXPANDR | 3 | R18. | Number of Trees Per Acre | 18 |
| EXPAREA | 1 | C7. | Area Expander | 11 |
| EXPREMEA | 1 | R28. | Remeasurement Expander | 11 |
| EXPSHUL | 5 | R4. | Expander for low shrubs | 21 |
| EXPSHUT | 4 | R5. | Expander for tall shrubs | 20 |
| EXPVOL | 2 | R29. | Volume Expander | 14 |
| FORTCUR | 2 | C4. | Current Forest Type | 14 |
| FORTORG | 2 | R23. | Original Forest Type | 14 |
| FORTPRJ | 2 | R20. | Projected Forest Type | 14 |
| GLUCUR | 1 | C2. | Ground Land Use Current | 10 |
| GLUORG | 1 | R6. | Ground Land Use Original | 10 |
| GLUPRJ | 1 | R7. | Ground Land Use Projected | 10 |
| GROPER | 1 | R15. | Remeasurement Period | 10 |
| GRSTKPC | 2 | R36. | Growing Stock Tree Stocking | 15 |
| GSCUVH | 2 | R31. | Growing Stock CuFt Volume Hardwoods | 15 |
| GSCUVS | 2 | R30. | Growing Stock CuFt Volume Softwoods | 15 |
| HTSI | 6 | R4. | Height Site Index Tree | 21 |
| IDENT | 1 | C6. | Case Identifier | 9 |
| IDENT | 2 | C6. | Case Identifier | 12 |
| IDENT | 3 | C6. | Case Identifier | 17 |
| IDENT | 4 | C6. | Case Identifier | 20 |
| IDENT | 5 | C6. | Case Identifier | 20 |
| IDENT | 6 | C6. | Case Identifier | 21 |
| IDENT | 7 | C6. | Case Identifier | 22 |
| LOGGR | 3 | R9. | Log Grade | 17 |
| MORTFAC | 3 | R20. | Mortality Factor | 18 |
| NATFOR | 2 | R6. | National Forest Code | 12 |
| NCSPGP | 3 | R21. | North Central Species Group | 19 |
| NETCUVL | 3 | R25. | Net Cubic Foot Volume in Tree | 19 |
| OWNGP | 2 | C3. | Ownership Group | 12 |
| PHOTAGE | 1 | R12. | Photo Age | 10 |
| PHYCLS | 2 | R12. | Physiographic Class | 13 |
| PICODE | 1 | R4. | Photo Interpretation Code | 9 |
| PISTRAT | 1 | R5. | Photo Stratification Code | 10 |
| POSITION | 2 | R10. | Position on Slope | 13 |
| PRJAGE | 2 | R26. | Projected Stand Age | 14 |
| PRJAREA1 | 1 | R20. | Projected Area Expander 1 | 11 |
| PRJAREA2 | 1 | R21. | Projected Area Expander 2 | 11 |
| PRJCRRAT | 3 | R12. | Projected Crown Ratio | 18 |
| PRJDATE | 1 | R14. | Projected Date | 10 |


(continued on next page)

INDEX


| Variable name | Record number | Variable number | Definition | Page |
|---|---|---|---|---|
| PRJEXPD | 3 | R19. | Projection Factor | 18 |
| PRJGSPC | 2 | R39. | Projected G.S. Tree Stocking | 16 |
| PRJHIST | 3 | R3. | Projected Tree History-Status | 17 |
| PRJHVOL | 2 | R33. | Projected G.S. CuFt Volume Hardwoods | 15 |
| PRJNCVL | 3 | R26. | Projected Net CuFt Volume in Tree | 19 |
| PRJOWN | 2 | R8. | Projected Owner Group | 13 |
| PRJRGPC | 2 | R40. | Projected Rough Tree Stocking | 16 |
| PRJRTPC | 2 | R41. | Projected Rotten Tree Stocking | 16 |
| PRJSTDH | 1 | R11. | Projected Stand History | 10 |
| PRJSVOL | 2 | R32. | Projected G.S. CuFt Volume Softwoods | 15 |
| PRJTRCLS | 3 | R16. | Projected Tree Class | 18 |
| PRJTRSZ | 3 | R23. | Projected Tree Size Class | 19 |
| PTCLS | 7 | R2. | Point Class | 22 |
| PTNUM | 7 | R1. | Point Number | 22 |
| PTOCC | 3 | R17. | Point Occupancy | 18 |
| RANDIST | 2 | R7. | Ranger District | 12 |
| RANGE | 1 | R26. | Range | 11 |
| ROTENPC | 2 | R38. | Rotten Tree Stocking | 16 |
| ROUGHPC | 2 | R37. | Rough Tree Stocking | 16 |
| SAMPKND | 1 | R2. | Sample Kind | 9 |
| SBLNUM | 5 | R1. | Shrub Number-Low | 20 |
| SBTNUM | 4 | R1. | Shrub Number-Tall | 20 |
| SECTION | 1 | R27. | Section | 11 |
| SEEDSOR | 2 | R14. | Seed source | 13 |
| SG01 | 2 | C8. | Presence or Absence of Jack Pine | 15 |
| SG02 | 2 | C9. | Presence or Absence of Red Pine | 15 |
| SG03 | 2 | C10. | Presence or Absence of White Pine | 15 |
| SG04 | 2 | C11. | Presence or Absence of Ponderosa Pine | 15 |
| SG05 | 2 | C12. | Presence or Absence of Loblolly Pine | 15 |
| SG06 | 2 | C13. | Presence or Absence of Shortleaf Pine | 15 |
| SG07 | 2 | C14. | Presence or Absence of Other Yellow Pines | 15 |
| SG08 | 2 | C15. | Presence or Absence of White Spruce | 15 |
| SG09 | 2 | C16. | Presence or Absence of Black Spruce | 15 |
| SG10 | 2 | C17. | Presence or Absence of Balsam Fir | 15 |
| SG11 | 2 | C18. | Presence or Absence of Hemlock | 15 |
| SG12 | 2 | C19. | Presence or Absence of Tamarack | 15 |
| SG13 | 2 | C20. | Presence or Absence of Bald Cypress | 15 |
| SG14 | 2 | C21. | Presence or Absence of E. Red Cedar | 15 |
| SG15 | 2 | C22. | Presence or Absence of N. white-cedar | 15 |
| SG16 | 2 | C23. | Presence or Absence of (not assigned) | 15 |
| SG17 | 2 | C24. | Presence or Absence of Other Softwoods | 15 |
| SG18 | 2 | C25. | Presence or Absence of Select White Oak | 15 |
| SG19 | 2 | C26. | Presence or Absence of Other White Oak | 15 |
| SG20 | 2 | C27. | Presence or Absence of Select Red Oak | 15 |
| SG21 | 2 | C28. | Presence or Absence of Other Red Oak | 16 |
| SG22 | 2 | C29. | Presence or Absence of Select Hickory | 16 |
| SG23 | 2 | C30. | Presence or Absence of Other Hickory | 16 |
| SG24 | 2 | C31. | Presence or Absence of Basswood | 16 |
| SG25 | 2 | C32. | Presence or Absence of Beech | 16 |
| SG26 | 2 | C33. | Presence or Absence of Yellow Birch | 16 |
| SG27 | 2 | C34. | Presence or Absence of Hard Maple | 16 |
| SG28 | 2 | C35. | Presence or Absence of Soft Maple | 16 |
| SG29 | 2 | C36. | Presence or Absence of Elm | 16 |
| SG30 | 2 | C37. | Presence or Absence of Black Ash | 16 |
| SG31 | 2 | C38. | Presence or Absence of White-Green Ash | 16 |
| SG32 | 2 | C39. | Presence or Absence of Sycamore | 16 |
| SG33 | 2 | C40. | Presence or Absence of Cottonwood | 16 |
| SG34 | 2 | C41. | Presence or Absence of Willow | 16 |
| SG35 | 2 | C42. | Presence or Absence of Hackberry | 16 |
| SG36 | 2 | C43. | Presence or Absence of Balsam Poplar | 16 |
| SG37 | 2 | C44. | Presence or Absence of Bigtooth Aspen | 16 |
| SG38 | 2 | C45. | Presence or Absence of Quaking Aspen | 16 |
| SG39 | 2 | C46. | Presence or Absence of Paper Birch | 16 |
| SG40 | 2 | C47. | Presence or Absence of River Birch | 16 |
| SG41 | 2 | C48. | Presence or Absence of Sweetgum | 16 |
| SG42 | 2 | C49. | Presence or Absence of Tupelo | 16 |


(Continued on next page)

| Variable name | Record number | Variable number | Definition | Page |
|---|---|---|---|---|
| SG43 | 2 | C50. | Presence or Absence of Black Cherry | 16 |
| SG44 | 2 | C51. | Presence or Absence of Black Walnut | 16 |
| SG45 | 2 | C52. | Presence or Absence of Butternut | 16 |
| SG46 | 2 | C53. | Presence or Absence of yellow Poplar | 16 |
| SG47 | 2 | C54. | Presence or Absence of (not assigned) | 16 |
| SG48 | 2 | C55. | Presence or Absence of (not assigned) | 16 |
| SG49 | 2 | C56. | Presence or Absence of Other Hardwoods | 16 |
| SG50 | 2 | C57. | Presence or Absence of Noncommercial | 16 |
| SHRUBSP | 4 | R2. | Shrub Species-Tall | 20 |
| SHUSPL | 5 | R2. | Shrub Species-Low | 20 |
| SISPP | 2 | R28. | Site Index Species | 14 |
| SITECLS | 2 | R27. | Site Class | 14 |
| SITEIND | 2 | C5. | Site Index | 14 |
| SLOPE | 2 | R11. | Slope Angle at Plot Center | 13 |
| SOILCLS | 1 | R19. | Soil Classification | 11 |
| SOUND | 3 | R8. | Soundness Code | 17 |
| SPECIES | 3 | R4. | Tree Species | 17 |
| SPPSI | 6 | R3. | Species Site Index Tree | 21 |
| STATE | 1 | R1. | State Code | 9 |
| STDAGE | 2 | R25. | Stand Age | 14 |
| STDAREA | 2 | R34. | Stand Area | 15 |
| STDHIST | 1 | R10. | Stand History | 10 |
| STDORG | 2 | R13. | Stand Origin | 13 |
| SURDATE | 1 | R13. | Survey Date | 10 |
| SZDNCUR | 2 | R21. | Current Stand Size-Density | 14 |
| SZDNORG | 2 | R24. | Original Stand Size-Density | 14 |
| SZDNPRJ | 2 | R22. | Projected Stand Size-Density | 14 |
| TOWNSHIP | 1 | R25. | Township | 11 |
| TRCLS | 3 | R15. | Tree Class | 18 |
| TRHIST | 3 | R2. | Tree History-Status | 17 |
| TRNUM | 3 | R1. | Tree Number | 17 |
| TRNUM | 6 | R1. | Tree Number | 21 |
| TRSIND | 3 | R24. | Tree Site Index | 19 |
| TRSZCLS | 3 | R22. | Tree Size Class | 19 |
| UNTCO | 1 | C1. | Unit and County | 9 |
| USEAREA | 1 | R17. | Use Area-Recreation | 11 |
| USEPOST | 1 | R18. | Use Posted-Recreation | 11 |
| USESET | 1 | R16. | Use Setting-Recreation | 10 |
| UTMEAST | 1 | R24. | UTM Easting To Nearest 1000 Meters | 11 |
| UTMNORTH | 1 | R23. | UTM Northing To Nearest 1000 Meters | 11 |
| UTMZONE | 1 | R22. | UTM Zone Code | 11 |
| UTREND | 1 | R9. | Use Trend | 10 |
| YRSADD | 6 | R6. | Years Added to DBH Age for Total Age | 21 |

## SECTION III.--STORED PROCEDURES

The Procedure File is a part of a SIR data base. "This file contains members which may be texts of commands, data or anything a user wishes to store with the data base. Typically, these texts will be frequently used SIR retrieval programs or other SIR command sequences.

"Each member in the Procedure File is identified by a member name and, optionally, a family name. Members are said to belong to a family and must have a unique name within the family." (Robinson et al. 1980).

We have created three families of procedures that we believe will be useful to users: COEF, LABEL, and VOLUME. Family COEF contains the coefficients needed to compute gross and net volumes using the equations presented by Hahn 1984, Smith and Weist 1982, Hahn 1976, Hahn 1975. Family LABEL contains procedures useful in making tables indexed by various data items. Family VOLUME contains the procedures for computing volumes in board feet, cubic feet, or green tons.

| Procedure | |
|---|---|
| COEF.BDFT:T | |
| COEF.CUFT:T | |
| COEF.CUSL:T | |
| COEF.GRTONS:T | |
| COEF.HEIGHT:T | |
| VOLUME.BDFT20:T | These volume procedures compute the gross and net |
| VOLUME.BDFT31:T | cubic or board foot volume for the indicated tree |
| VOLUME.CUFT20:T | class from the tree d.b.h., top diameter outside bark |
| VOLUME.CUFT30:T | and species group. They are implemented by |
| VOLUME.CUFT40:T | CALL VOLUME.xxxxxx(Dbh,tdob,sp.gp.) |
| VOLUME.GRTONS:T | Computes biomass in green tons. |
| VOLUME.HEIGHT:T | Computes total or merchantable height. |

| Procedure | Description |
|---|---|
| LABEL.BACLASS:T | Computes an index and labels in variable BACLS for stand basal area using the variable BACUR. |
| LABEL.CONUNS:T | Computes an index and labels in variable CONUN for conifer understory using the variable CUNCOND. |
| LABEL.DBHCLS:T | Computes an index and labels in variables DIACLS1, DIACLS5, and DIACLS9 for 2-inch diameter class using the variable DBHCUR. |
| LABEL.DEATH:T | Computes an index and labels in variable DEATH for cause of death using the variable DAMDETH. |
| LABEL.LANDCLS:T | Computes an index and labels in variable LANDUSE for ground land use using the variables GLUCUR and FORTCUR. |
| LABEL.LIVE:T | Computes an index and labels in variable LIVE for tree history of live trees using the variable TRHISTCU. |
| LABEL.LOGGRD:T | Computes an index and labels in variable LOGGRD for tree log grade using the variable LOGGR. |
| LABEL.OWNER:T | Computes an index and labels in variable OWNER for stand owner class using the variable OWNGP. |
| LABEL.SICLS:T | Computes an index and labels in variable SICLS for stand site index using the variable SITEIND. |
| LABEL.SPECIES:T | Computes an index and labels in variables SPECIEC and SPECIESN for individual tree species of commercial and noncommercial species, respectively, using the variable SPECIES. |
| LABEL.STCLGR:T | Computes an index and labels in variable BACLS for stand basal area using the variable BACUR. |
| LABEL.STDAGE:T | Computes an index and labels in variable STDAGECL for stand age using the variable STDAGE. |
| LABEL.STDVOL:T | Computes an index and labels in variable STDVOLCL for stand volume class using the variables GSCUVS and GSCUVH. |
| LABEL.SWDHWD:T | Computes an index and labels in variable SWDHWD for major tree species group using the variable SPECIES. |
| LABEL.TRCLASS:T | Computes an index and labels in variable TRCLASS for tree class using the variable TRCLS. |

| | |
|---|---|
| LABEL.TYPESZ:T | Computes an index and labels in variables FORTYPE and SZCLS for forest type and stand size class, respectively, using the variables FORTCUR and SZDNCUR. |
| LABEL.U1COUNTY:T<br>LABEL.U2COUNTY:T<br>LABEL.U3COUNTY:T<br>LABEL.U4COUNTY:T<br>LABEL.U5COUNTY:T | These procedures compute an index and labels in variable COUNTY for county code using the variable UNTCO. |

## SECTION IV.--EXAMPLE RETRIEVALS

The following is a simple retrieval that will produce a table of area of commercial forest land by ownership group and forest type for a five-county area in central Wisconsin. The submit file shown below has two main parts, the Cyber control statements and the SIR retrieval statements. The Cyber control statements are the first part of the file, up through "/EOR" statement. These statements identify the user number, set time limits, execute SIR, save output files, and send the output files to the printer. This part of the submit file will be basically the same for all retrievals. For information on these commands, see the University of Minnesota Computer Center User's Manual. The remainder of the submit file contains SIR retrieval statements that are used to retrieve the needed data and construct the output table. This part of the submit file will be different for every retrieval. For information on these commands, see the SIR Version 2 User's Manual and SIR/DBMS TABLE Procedure Reference Guide.

```
*******************************************************************************

/JOB
/NOSEQ
MHH,T20.
/USER
FETCH,SIR.
SIR.
RETAIN,EXAM1TB.
GOFO,COST.
EXIT.
COST.
DAYFILE.
RETAIN,OUTPUT=EXAM1OT.
ROUTE,OUTPUT,DC=PR,TID=BC,BIN=R69,EC=A9.
ROUTE,EXAM1OT,DC=PR,TID=BC,BIN=R69,EC=A9.
/EOR
OLD FILE        WISSOU
PASSWORD        WISSOU
SECURITY        WISSOU,WISSOU
PAGESIZE        63,89
RETRIEVAL
PROCESS CASE    LIST = 300000 THRU 399999
INCLUDE         FORTYPE,OWNGP,EXPAREA


.  IF     (GLUCUR GT 22) NEXT CASE
.  IFNOT  (UNTCO EQ 301 OR 308 OR 310 OR 311 OR 312 ) NEXT CASE
.  PROCESS REC 2
.    MOVE VARS OWNGP,EXPAREA
.    CALL LABEL.TYPESZ
.  END PROCESS REC
.  PERFORM PROCS
END PROCESS CASE
TABLE FORTYPE.T, OWNGP.T BY EXPAREA.O/
      FILENAME=EXAM1TB/
      PAGELENGTH=63/PAGEWIDTH=89/HEADERWIDTH=12/
      STUBTITLE = 'Forest type'/
      WAFERTITLE =
    'Example 1.--Area of commercial forest land by ownership'
    'and forest type, Adams, Marquette, Portage, Waupaca, and'
    'Waushara counties, Wisconsin, 1983.'
    'North Central Forest Experiment Station FI&A Data Base.'
    '     '
    '(In acres)'/
      NOOVERPRINTING/NOBOLDPRINTING
END RETRIEVAL
FINISH
```

The following table was produced by this retrieval and retained in the file EXAM1TB.

**********

/DBMS Table Procedure Page 1

Example 1.--Area of commercial forest land by ownership and forest type, Adams, Marquette, Portage, Waupaca, and Waushara counties, Wisconsin, 1983.

North Central Forest Experiment Station FI&A Data Base.

(In acres)

| Forest type | TOTAL | National forest | Other public | Private |
|---|---|---|---|---|
| | Area expander | | Area expander | |
| TOTAL..... | 821100 | 0 | 31600 | 789500 |
| Jack pine..... | 51500 | 0 | 0 | 51500 |
| Red pine..... | 85700 | 0 | 0 | 85700 |
| White pine..... | 25100 | 0 | 2300 | 22800 |
| Balsam fir..... | 7200 | 0 | 0 | 7200 |
| White spruce..... | 0 | 0 | 0 | 0 |
| Black spruce..... | 0 | 0 | 0 | 0 |
| Northern white-cedar | 13900 | 0 | 0 | 13900 |
| Tamarack..... | 13700 | 0 | 0 | 13700 |
| Oak-hickory..... | 290300 | 0 | 11500 | 278800 |
| Elm-ash-soft maple.. | 100400 | 0 | 2200 | 98200 |
| Maple-birch..... | 103400 | 0 | 4400 | 99000 |
| Aspen..... | 91100 | 0 | 11200 | 79900 |
| Paper birch..... | 18300 | 0 | 0 | 18300 |
| Exotic..... | 2200 | 0 | 0 | 2200 |
| Nonstocked..... | 18300 | 0 | 0 | 18300 |

The following listing was produced in example 1 and retained in the file EXAM1OT.

```
*******************************************************************************

 R / D B M S  2 . 1 . 1       (SIR/DBMS 2.1.1)    01/12/84  16:15:52  PAGE     1

          OLD FILE          WISSOU
          PASSWORD       XXXXXXXXXXXXXXX
          SECURITY       XXXXXXXXXXXXXXX
          PAGESIZE          63,89
    5     RETRIEVAL
          PROCESS CASE      LIST = 300000 THRU 399999
          INCLUDE           FORTYPE,OWNGP,EXPAREA
          .  IF     (GLUCUR GT 22) NEXT CASE
          .  IFNOT  (UNTCO EQ 301 OR 308 OR 310 OR 311 OR 312 ) NEXT CASE
   10     .  PROCESS REC 2
          .    MOVE VARS OWNGP,EXPAREA
          .    CALL LABEL.TYPESZ
1.10      .    RECODE       FORTYPE=FORTCUR(1=1)(2=2)(3=3)(6=14)(13=4)(12=6)
1.20                                       (16=5)(14=7)(15=8)(50=9)(70=10)
1.30                                       (80=11)(91=12)(92=13)
1.40      .    IF           (SZDNCUR EQ 40) FORTYPE = 15
1.50      .    COMPUTE      STDSZCLS = TRUNC(SZDNCUR/10)
1.60      .    VAR RANGES   STDSZCLS(1,4)/
1.70                        FORTYPE(1,15)
1.80      .    VAR LABELS   STDSZCLS Stand size class/
1.90                        FORTYPE  Forest type
1.100     .    VALUE LABELS STDSZCLS (1) Sawtimber
1.110                                 (2) Poletimber
1.120                                 (3) Sapling & seedling
1.130                                 (4) Nonstocked/
1.140                       FORTYPE  (1) Jack pine
1.150                                (2) Red pine
1.160                                (3) White pine
1.170                                (4) Balsam fir
1.180                                (5) White spruce
1.190                                (6) Black spruce
1.200                                (7) Northern white-cedar
1.210                                (8) Tamarack
1.220                                (9) Oak-hickory
1.230                                (10) Elm-ash-soft maple
1.240                                (11) Maple-birch
1.250                                (12) Aspen
1.260                                (13) Paper birch
1.270                                (14) Exotic
1.280                                (15) Nonstocked
          .  END PROCESS REC
          .  PERFORM PROCS
   15     END PROCESS CASE
          TABLE FORTYPE.T, OWNGP.T BY EXPAREA.O/

               FILENAME=EXAM1TB/
               PAGELENGTH=63/PAGEWIDTH=89/HEADERWIDTH=12/
               STUBTITLE = 'Forest type'/
   20          WAFERTITLE =
              'Example 1.--Area of commercial forest land by ownership'
              'and forest type, Adams, Marquette, Portage, Waupaca, and'
              'Waushara counties, Wisconsin, 1983.'
              'North Central Forest Experiment Station FI&A Data Base.'
   25         '     '
              '(In acres)'/
                NOOVERPRINTING/NOBOLDPRINTING
          END RETRIEVAL
          FINISH
```

```
FULL  DAYFILE.  84/01/12. 16.19.19.* AKZIBDK* PAGE     1

 AKZIBDK. 84/01/12.UOFM CYBER A (01/02-BO).
16.15.44.MHH,T20.
16.15.44.$USER(WXL6007,)
16.15.44. FAMILY PARAMETER MISSING.  FAMILY CA ASSUMED.
16.15.46.FETCH,SIR.
16.15.52.SIR.
16.15.53.*** REMARK ***  BEGIN SIR/DBMS 2.1.1 RUN.
16.15.55.*** REMARK ***  TASK TABLE SPACE USED:    126
16.15.55.*** REMARK ***  TOTAL CPU TIME USED (MS): 37
16.15.56.*** REMARK ***   START RETRIEVAL TRANSLATION.
16.16.05.*** REMARK ***   START RETRIEVAL EXECUTION.
16.19.17.*** REMARK ***   RETRIEVAL COMPLETE.
16.19.18.*** REMARK ***  TASK TABLE SPACE USED:    4355
16.19.18.*** REMARK ***  TOTAL CPU TIME USED (MS): 7845
16.19.18.*** REMARK ***  END SIR RUN:          0
16.19.18.*** REMARK ***  TOTAL CPU TIME (MS): 7855
16.19.18.*** REMARK ***  TABLE SPACE    USED: 4355
16.19.18.*** REMARK ***  FIELD LENGTH   USED: 36800
16.19.19.RETAIN,EXAM1TB.
16.19.19. EXAM1TB PURGED.
16.19.19. PERMANENT FILE EXAM1TB, 7 PRUS.
16.19.19.GOFO,COST.
16.19.19.COST.
16.19.19. ACCUMULATED JOB COST - UNIVERSITY RATE.
16.19.19. CP       7.906 SEC.
16.19.19. MS      57.732 KUN.
16.19.19. PF        .108 KUN.
16.19.19. CM      69.470 KWD.
16.19.19. SR      21.555 UNS.
16.19.19. CURRENT JOB COST =     $ 3.23
16.19.19.DAYFILE.

*******************************************************************************
```

Example 1.

This is a simple retrieval that produces a commonly requested table for a specific area. The retrieval uses the PROCESS CASE command to restrict the retrieval to the desired forest survey unit. An IF statement is used to further restrict the retrieval to only commercial forest land and an IFNOT statement is used to restrict it to the desired five-county area. Becasuse the retrieval requires only data stored in record type 2 , only that record is accessed. Two of the variables required to construct the table (OWNGP and AREAEXP) are stored as needed in the data base and can be moved to the summary record with a MOVE VARS command. A stored procedure TYPESZ is called to compute FORTYPE from FORTCUR to reduce the space needed to produce the output table. The cost shown for this example is based on daytime rates for university or affiliated users. Higher rates are charged to private and corporate users; lower rates are charged for jobs submitted to be run in the evening or overnight.

Two more example retrievals follow. Although these retrievals are more complicated, they are basically extensions of the first example. Only the SIR retrieval statements section of the submit file are shown for these examples because the Cyber control statements do not differ.

Example 2.

This retrieval produces two different outputs, a table and a data file, for a single county. The table presents number of ground plots, area of commercial forest land, and average softwood and hardwood volume per acre by forest type. The data file contains two types of records: (1) a plot record that is identified by a 1 in the first column and (2) tree records that have a 2 in column one.

```
*******************************************************************************

OLD FILE          WISSOU
PASSWORD          WISSOU
SECURITY          WISSOU,WISSOU
PAGESIZE          63,89
RETRIEVAL
PROCESS CASE      LIST = 300000 THRU 399999
INCLUDE           IDENT,FORTCUR,EXPAREA,OWNGP,SITEIND,
                  TRNUM,DBHCUR,TRHIST,NCSPGP,EXPANDR,GSCUVS,GSCUVH
.  IFNOT  (UNTCO EQ 302) NEXT CASE
.  IF     (GLUCUR GT 22) NEXT CASE
.  PROCESS REC 2
.     MOVE VARS IDENT,FORTCUR,OWNGP,SITEIND,GSCUVS,GSCUVH,EXPAREA
COMMENT ========== COMPUTE FOREST TYPE ====================
.     CALL LABEL.TYPESZ
COMMENT ========== ROUND ACRES FOR TABLE ====================
.     COMPUTE    MACRES = RND(EXPAREA/100)/10
.     VAR LABEL  MACRES Area (Thousand acres)/
COMMENT ========== GENERATE A PLOT RECORD ====================
.     WRITE       '1 ' IDENT (I6), UNTCO (I3), FORTCUR (I4), EXPAREA (I7),
                  OWNGP (I3), SITEIND (I5)
.  END PROCESS REC
.  PROCESS REC   3
.     MOVE VARS    IDENT,TRNUM,DBHCUR,TRHIST,NCSPGP,EXPANDR
COMMENT ========== GENERATE A TREE RECORD ====================
.     WRITE       '2 ' IDENT (I6), TRNUM (I5), DBHCUR (F5.1), TRHIST (I3),
                  NCSPGP (I3), EXPANDR (I3)
.  END PROCESS REC
.  PERFORM PROCS
END PROCESS CASE
TABLE FORTYPE THEN TOTFT, N THEN MACRES.O THEN (GSCUVS THEN GSCUVH) BY MEAN/
      FILENAME=EXAM2TB/
      PRINTFORMAT = N(C,0) MACRES(C,1) CSUVS(C,0) GSCUVH(C,0)/
      TOTAL= N 'Number of plots' TOTFT 'All types'/
      MEAN = MEAN '(cubic feet per acre)'/
      PAGETITLE = 'Example 2.'/
          PAGELENGTH = 63/
      PAGEWIDTH = 89/
      HEADERDIVIDER = ' '/
      STUBDIVIDER = '-'/
      STUBFILLER = ' '/
      HEADERWIDTH = 12/
      STUBTITLE = 'Forest type'/
      WAFERTITLE =
      'Example 2.--Number of plots, area of commercial forest land,'
      'and average softwood and hardwood volume per acre,'
      'Chippewa County, Wisconsin, 1983'/
      NOLEFTBORDER/NORIGHTBORDER/
      NOZEROS/NOEMPTYROWS/
      NOOVERPRINTING/NOBOLDPRINTING
END RETRIEVAL
FINISH
```

The following table was produced by this retrieval and retained in the file EXAM2TB.

**********

Example 2. Page 1

Example 2.--Number of plots, area of commercial forest land, and average softwood and hardwood volume per acre, Chippewa County, Wisconsin, 1983

| Forest type | Number of plots | Area (Thousand acres) | Growing stock cubic foot volume softwood (cubic feet per acre) | Growing stock cubic foot volume hardwood (cubic feet per acre) |
|---|---|---|---|---|
| Jack pine | 1 | 2.5 | 631 | - |
| White pine | 2 | 5.0 | 1496 | 148 |
| Black spruce | 1 | 2.4 | 72 | - |
| Tamarack | 3 | 7.3 | 520 | 263 |
| Oak-hickory | 11 | 27.1 | 87 | 1,071 |
| Elm-ash-soft maple | 16 | 39.4 | 15 | 596 |
| Maple-birch | 32 | 77.8 | 31 | 938 |
| Aspen | 19 | 47.0 | 34 | 1,040 |
| Paper birch | 6 | 14.7 | 114 | 749 |
| Nonstocked | 1 | 2.5 | - | 133 |
| All types | 92 | 225.7 | 95 | 835 |

The following listing was produced in example 2 and retained in the file EXAM2OT. The data file was written by the two WRITE statements and is part of this output file. Only a portion of the data file is shown.

**********

```
S I R / D B M S  2 . 1 . 1          (SIR/DBMS 2.1.1)        01/23/84  16:25:07  PAGE     1

           OLD FILE          WISSOU
           PASSWORD       XXXXXXXXXXXXXXX
           SECURITY       XXXXXXXXXXXXXXX
           PAGESIZE          63,89
    5      RETRIEVAL
           PROCESS CASE      LIST = 300000 THRU 399999
           INCLUDE           IDENT,FORTCUR,EXPAREA,OWNGP,SITEIND,
                             TRNUM,DBHCUR,TRHIST,NCSPGP,EXPANDR,GSCUVS,GSCUVH
           .  IFNOT  (UNTCO EQ 302) NEXT CASE
   10      .  IF     (GLUCUR GT 22) NEXT CASE
           .  PROCESS REC 2
           .    MOVE VARS IDENT,FORTCUR,OWNGP,SITEIND,GSCUVS,GSCUVH,EXPAREA
           COMMENT ========== COMPUTE FOREST TYPE ====================
           .    CALL LABEL.TYPESZ
               .    CALL LABEL.TYPESZ
  1.10 .   RECODE       FORTYPE=FORTCUR(1=1)(2=2)(3=3)(6=14)(13=4)(12=6)
  1.20                                 (16=5)(14=7)(15=8)(50=9)(70=10)
  1.30                                 (80=11)(91=12)(92=13)
  1.40 .   IF           (SZDNCUR EQ 40) FORTYPE = 15
  1.50 .   COMPUTE      STDSZCLS = TRUNC(SZDNCUR/10)
  1.60 .   VAR RANGES   STDSZCLS(1,4)/
  1.70                  FORTYPE(1,15)
  1.80 .   VAR LABELS   STDSZCLS Stand size class/
  1.90                  FORTYPE  Forest type
  1.100 .  VALUE LABELS STDSZCLS (1) Sawtimber
  1.110                          (2) Poletimber
```

```
1.120                          (3) Sapling & seedling
1.130                          (4) Nonstocked/
1.140                FORTYPE   (1) Jack pine
1.150                          (2) Red pine
1.160                          (3) White pine
1.170                          (4) Balsam fir
1.180                          (5) White spruce
1.190                          (6) Black spruce
1.200                          (7) Northern white-cedar
1.210                          (8) Tamarack
1.220                          (9) Oak-hickory
1.230                          (10) Elm-ash-soft maple
1.240                          (11) Maple-birch
1.250                          (12) Aspen
1.260                          (13) Paper birch
1.270                          (14) Exotic
1.280                          (15) Nonstocked

  15      COMMENT ========== ROUND ACRES FOR TABLE ====================
          .    COMPUTE    MACRES = RND(EXPAREA/100)/10
          .    VAR LABEL  MACRES Area (Thousand acres)/
          COMMENT ========== GENERATE A PLOT RECORD ====================
          .    WRITE       '1 ' IDENT (I6), UNTCO (I3), FORTCUR (I4), EXPAREA (I7),
  20                       OWNGP (I3), SITEIND (I5)
          .  END PROCESS REC
          .  PROCESS REC   3
          .    MOVE VARS    IDENT,TRNUM,DBHCUR,TRHIST,NCSPGP,EXPANDR
          COMMENT ========== GENERATE A TREE RECORD ====================
  25      .    WRITE       '2 ' IDENT (I6), TRNUM (I5), DBHCUR (F5.1), TRHIST (I3),
                            NCSPGP (I3), EXPANDR (I3)
          .  END PROCESS REC
          .  PERFORM PROCS
          END PROCESS CASE

  30      TABLE FORTYPE THEN TOTFT, N THEN MACRES.O THEN (GSCUVS THEN GSCUVH) BY MEAN/
                FILENAME=EXAM2TB/
                PRINTFORMAT = N(C,0) MACRES(C,1) CSUVS(C,0) GSCUVH(C,0)/
                TOTAL= N 'Number of plots' TOTFT 'All types'/
                MEAN = MEAN '(cubic feet per acre)'/
  35            PAGETITLE = 'Example 2.'/
                PAGELENGTH = 63/
                PAGEWIDTH = 89/
                HEADERDIVIDER = ' '/
                STUBDIVIDER = '-'/
  40            STUBFILLER = ' '/
                HEADERWIDTH = 12/
                STUBTITLE = 'Forest type'/
                WAFERTITLE =
                'Example 2.--Number of plots, area of commercial forest land,'
  45            'and average softwood and hardwood volume per acre,'
                'Chippewa County, Wisconsin, 1983'/
                NOLEFTBORDER/NORIGHTBORDER/
                NOZEROS/NOEMPTYROWS/
                NOOVERPRINTING/NOBOLDPRINTING
  50      END RETRIEVAL
1 300197302  50   2500  4   84
2 300197  101  6.0  1 20 19
2 300197  102  4.3  1 20100
2 300197  201 13.3  1 38  4
2 300197  202  7.3  1 21 13
2 300197  203  6.0  1 18 19
2 300197  301 13.9  1 38  4
2 300197  302  9.1  1 21  8
2 300197  401  1.5  2 50  0
2 300197  402  1.7  2 50  0
          .
          .
          .
1 354220302  91   2400  4   64
```

```
2 354220   101   0.0 31 38100
2 354220   102   0.0 31 38100
2 354220   201   1.0 31 38100
2 354220   301   0.0 31 38100
2 354220   302   0.0 31 38100
2 354220   303   0.0 31 38100
2 354220   401   0.0 31 38   0
2 354220   402   2.2 31 38   0
2 354220   601   1.6 61 38   0
2 354220   701   0.0 31 38   0
2 354220   702   0.0 31 38   0
2 354220   703   0.0 31 38   0
2 354220   704   0.0 31 38   0
2 354220   705   0.0 31 38   0
2 354220   706   0.0 31 38   0
2 354220   707   0.0 31 38   0
2 354220   801   1.7 61 38   0
2 354220   901   0.0 31 39   0
2 354220   902   0.0 31 28   0
2 354220   903   0.0 31 28   0
            FINISH
```

**********

This example was run at normal rates, used 10.995 seconds of central processing time, and cost $4.21 at university rates.

Example 3.

The following retrieval extracts the data for all plots located in a geographic area defined as a circle or a convex polygon, using Universal Transverse Mercator (UTM) coordinates. This retrieval also demonstrates the use of arrays, a FOR loop, and a WHILE loop to extract data for a pentagon-shaped area in the Upper Peninsula of Michigan. Only the retrieval is shown; any appropriate tables could be constructed using the selected plots and the previously described TABLE procedures.

```
**************************************************************************
OLD FILE            MICHUP
PASSWORD            MICHUP
SECURITY            MICHUP,MICHUP
PAGESIZE            58,136
RETRIEVAL
CALL COEF.HEIGHT
CALL COEF.CUFT
. EXCLUDE           C2A1 TO C2A50,C2B1 TO C2B50,C2C1 TO C2C50,C3C1 TO C3C50,
                    C4C1 TO C4C50,HTA1 TO HTA50,HTB1 TO HTB50,HTC1 TO HTC50,
                    HTD1 TO HTD50,HTE1 TO HTE50,HTF1 TO HTF50
COMMENT Procedure to determine if a given UTM location is within
            a specified circle or convex polygon. The following
            variables are used.
            NCORNER = The number of sides(corners/vertices) of the
                      polygon. If NCORNER = 1 a circle is assumed.
                      If NCORNER = 0 a county retrieval is assumed.
```

```
          NZONE    = The UTM zone for the supplied coordinates.
          XCORD1 TO XCORD10 = Up to 10 values for the EASTING

                    coordinate of each of the verticies.
          YCORD1 TO YCORD10 = Up to 10 values for the NORTHING

                    coordinate of each of the verticies.
          If NCORNER = 1 XCORD1, YCORD1 is the coordinate of

                    the center of the circle.
             NCORNER = 0 XCORD contains Unit,County(UNTCO) codes

                    for counties to be retrieved and RADIUS

                    contains the number of counties to be retrieved.

          RADIUS = Radius of the circle in kilometers for a circle

                   retrieval or the number of counties for a county

                   retrieval.

.  SET              NCORNER,RADIUS,XCORD1 TO XCORD10,YCORD1 TO YCORD10,
                    XMAX,XMIN,YMAX,YMIN,XT1 TO XT3,YT1 TO YT3,A1 TO A3,
                    B1 TO B3,V1 TO V3(0)
COMMENT Enter data to define search area.
.  COMPUTE          NCORNER = 5;
                    RADIUS = 80;
                    NZONE = 16
COMMENT Enter coordinates of verticies XCORD are Easting and
        YCORD are northing.
.  COMPUTE          XCORD1 =  323;YCORD1 = 5194;
                    XCORD2 =  350;YCORD2 = 5160;
                    XCORD3 =  330;YCORD3 = 5101;
                    XCORD4 =  200;YCORD4 = 5140;
                    XCORD5 =  290;YCORD5 = 5180;
                    XCORD6 =    0;YCORD6 =    0;
                    XCORD7 =    0;YCORD7 =    0;
                    XCORD8 =    0;YCORD8 =    0;
                    XCORD9 =    0;YCORD9 =    0;
                    XCORD10=    0;YCORD10=    0
.  IFTHEN           (NCORNER GE 3)
COMMENT Define square containing subject polygon.
.    COMPUTE        XMAX = XCORD1; YMAX = YCORD1;
                    XMIN = XCORD1; YMIN = YCORD1
.    FOR            I = 2,NCORNER
.      COMPUTE      XTEST = XCORD1 TO XCORD10(I);
                    YTEST = YCORD1 TO YCORD10(I)
.      IF           (XTEST GT XMAX) XMAX = XTEST
.      IF           (XTEST LT XMIN) XMIN = XTEST
.      IF           (YTEST GT YMAX) YMAX = YTEST
.      IF           (YTEST LT YMIN) YMIN = YTEST
.    END FOR
.  ELSE
.  ENDIF
SET                 NPLTS,NTRS,NSUM(0)
PROCESS CASES       SAMPLE=.20,31
COMMENT Select commercial forest plots only.
.  IFNOT            (GLUCUR LE 29) NEXT CASE
.  PROCESS REC      1
.    COMPUTE        UTME=UTMEAST;UTMN=UTMNORTH
.    MOVE VARS      UNTCO
COMMENT OUT = 1 is true; 0 is false.
.    IFTHEN         (NCORNER EQ 0)
```

```
COMMENT Search is a county retrieval.
.      COMPUTE      OUT = 1
.      FOR          I = 1,RADIUS
.        IF         (UNTCO EQ XCORD1 TO XCORD10(I)) OUT = 0
.      END FOR
.    ELSEIF         (NCORNER EQ 1)
COMMENT Search area is a circle.
    Correct easting for zone change. 460 kilometers used for
       northern Michigan and Wisconsin.
.      IFTHEN       (UTMZONE NE NZONE)
.        IFTHEN     (UTMZONE LT NZONE)
.          COMPUTE UTME = UTME - 460
.        ELSE
.          COMPUTE UTME = UTME + 460
.        ENDIF
.      ELSE
.      ENDIF
.      COMPUTE      OUT = 1
.      COMPUTE      DISTANCE = (UTME-XCORD1)**2 + (UTMN-YCORD1)**2
.      IF           ((RADIUS - DISTANCE) GE 0.0) OUT = 0
.    ELSE
COMMENT Search area is a polygon.
       Correct easting for zone change. 460 kilometers used for
       northern Michigan and Wisconsin.
.      IFTHEN       (UTMZONE NE NZONE)
.        IFTHEN     (UTMZONE LT NZONE)
.          COMPUTE UTME = UTME - 460
.        ELSE
.          COMPUTE UTME = UTME + 460
.        ENDIF
.      ELSE
.      ENDIF
.      IFTHEN       (UTME GT XMAX OR UTME LT XMIN OR
                     UTMN GT YMAX OR UTMN LT YMIN)
.        COMPUTE    OUT = 1
.      ELSE
.        COMPUTE    OUT = 1;
                    ICOUNT = 0;
                    NCORNERT = NCORNER-2
.        WHILE      (ICOUNT LT NCORNERT AND OUT EQ 1)
.          COMPUTE ICOUNT = ICOUNT + 1;
                    ICOUN1 = ICOUNT + 1;
                    ICOUN2 = ICOUNT + 2;
                    XT1 = XCORD1;
                    XT2 = XCORD1 TO XCORD10(ICOUN1);
                    XT3 = XCORD1 TO XCORD10(ICOUN2);
                    YT1 = YCORD1;
                    YT2 = YCORD1 TO YCORD10(ICOUN1);
                    YT3 = YCORD1 TO YCORD10(ICOUN2);
                    A1 = (YT3-YT2) / (XT2*YT3 - XT3*YT2);
                    A2 = (YT3-YT1) / (XT1*YT3 - XT3*YT1);
                    A3 = (YT2-YT1) / (XT1*YT2 - XT2*YT1);
                        B1 = (XT3-XT2) / (XT2*YT3 - XT3*YT2);
                    B2 = (XT3-XT1) / (XT1*YT3 - XT3*YT1);
                    B3 = (XT2-XT1) / (XT1*YT2 - XT2*YT1);
                    V1 = (A1*UTME + B1*UTMN - 1.0) *
                         (A1*XT1 + B1*YT1 - 1.0);
                    V2 = (A2*UTME + B2*UTMN - 1.0) *
                         (A2*XT2 + B2*YT2 - 1.0);
                    V3 = (A3*UTME + B3*UTMN - 1.0) *
                        (A3*XT3 + B3*YT3 - 1.0)
.          IF       (V1 GE 0.0 AND V2 GE 0.0 AND V3 GE 0.0) OUT = 0
.        END WHILE
.      ENDIF
.    ENDIF
COMMENT If plot is not in specified region reject case.
.    IFNOT          (OUT EQ 0) NEXT CASE
.  END PROCESS REC
.  PROCESS REC      2
.    MOVE VARS      BACUR,SITEIND
```

```
.  END PROCESS REC
.  PROCESS REC      3
.    COMPUTE        STATUS = MOD(TRHIST,10)
COMMENT Select live trees only
.    IFNOT          (STATUS LE 2) NEXT REC
COMMENT Select pole sized and larger trees only
.    IFNOT          (TRSZCLS GE 2) NEXT REC
.    COMPUTE        NTRS = NTRS + 1
.    IFTHEN         (TRCLS LE 20)
.      COMPUTE      CUFTORG = 0.0
CALL VOLUME.CUFT20(DBHORG,4.0,NCSPGP)
.      COMPUTE      CUFTORG = CUFTNET
CALL VOLUME.CUFT20(DBHCUR,4.0,NCSPGP)
COMMENT Compute net growth
.      COMPUTE      NETGRO = ((CUFTNET-CUFTORG)*EXPANDR-CUFTNET*MORTFAC)*EXPAREA
.    ELSEIF         (TRCLS EQ 40)
CALL VOLUME.CUFT40(DBHCUR,4.0,NCSPGP)
.    ELSE
CALL VOLUME.CUFT30(DBHCUR,4.0,NCSPGP)
.    ENDIF
COMMENT Compute number of trees
.    COMPUTE        NUMTRS = EXPANDR*EXPAREA
COMMENT Compute cubic foot volume
.    COMPUTE        NETVOL = CUFTNET*EXPANDR*EXPAREA
.    COMPUTE        NSUM = NSUM + 1
.    PERFORM PROCS
.  END PROCESS REC
END PROCESS CASES
BLANK LINES     10
COMMENT List records processed in retrieval
WRITE 10X'PLOTS PROCESSED = ',NPLTS/10X,'TREES PROCESSED = ',NTRS
     /10X'SUMMARY RECORDS = ',NSUM
```

## APPENDIX A.--DEFINITION OF CODES

Codes for variable C1 UNTCO

State = Michigan  State code = 26
Data base number and name  (I) MICHUP

| Code | County name | Code | County name | Code | County name | Code | County name |
|---|---|---|---|---|---|---|---|
| Eastern Upper Peninsula | | | | | | | |
| (101) | Alger | (102) | Chippewa | (103) | Delta | (104) | Luce |
| (105) | Mackinac | (106) | Menominee | (107) | Schoolcraft | | |
| Western Upper Peninsula | | | | | | | |
| (201) | Baraga | (202) | Dickinson | (203) | Gogebic | (204) | Houghton |
| (205) | Iron | (206) | Keweenaw | (207) | Marquette | (208) | Ontonagon |

Data base number and name  (II) MICHLP

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Northern Lower Peninsula | | | | | | | |
| (301) | Alcona | (302) | Alpena | (303) | Antrim | (304) | Arenac |
| (305) | Bay | (306) | Benzie | (307) | Charlevoix | (308) | Cheboygan |
| (309) | Clare | (310) | Crawford | (311) | Emmet | (312) | Gladwin |
| (313) | Grand Traverse | (314) | Iosco | (315) | Isabella | (316) | Kalkaska |
| (317) | Lake | (318) | Leelanau | (319) | Manistee | (320) | Mason |
| (321) | Mecosta | (322) | Midland | (323) | Missaukee | (324) | Montmorency |
| (325) | Newaygo | (326) | Oceana | (327) | Ogemaw | (328) | Osceola |
| (329) | Oscoda | (330) | Otsego | (331) | Presque Isle | (332) | Roscommon |
| (333) | Wexford | | | | | | |
| Southern Lower Peninsula | | | | | | | |
| (401) | Allegan | (402) | Barry | (403) | Berrien | (404) | Branch |
| (405) | Calhoun | (406) | Cass | (407) | Clinton | (408) | Eaton |
| (409) | Genesee | (410) | Gratiot | (411) | Hillsdale | (412) | Huron |
| (413) | Ingham | (414) | Ionia | (415) | Jackson | (416) | Kalamazoo |
| (417) | Kent | (418) | Lapeer | (419) | Lenawee | (420) | Livingston |
| (421) | Macomb | (422) | Monroe | (423) | Montcalm | (424) | Muskegon |
| (425) | Oakland | (426) | Ottawa | (427) | Saginaw | (428) | St. Clair |
| (429) | St. Joseph | (430) | Sanilac | (431) | Shiawassee | (432) | Tuscola |
| (433) | Van Buren | (434) | Washtenaw | (435) | Wayne | | |

State = Wisconsin  State code = 55
Data base number and name  (III) WISNOR

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Northeast Unit | | | | | | | |
| (101) | Florence | (102) | Forest | (103) | Langlade | (104) | Lincoln |
| (105) | Menominee | (106) | Marinatte | (107) | Oconto | (108) | Oneida |
| (109) | Shawano | (110) | Vilas | | | | |
| Northwest Unit | | | | | | | |
| (201) | Ashland | (202) | Barron | (203) | Bayfield | (204) | Burnett |
| (205) | Douglas | (206) | Iron | (207) | Polk | (208) | Price |
| (209) | Rusk | (210) | Sawyer | (211) | Taylor | (212) | Washburn |

| Code | County name | Code | County name | Code | County name | Code | County name |
|---|---|---|---|---|---|---|---|

Data base number and name (IV) WISSOU

Central Unit

| | | | |
|---|---|---|---|
| (301) Adams | (302) Chippewa | (303) Clark | (304) Eau Claire |
| (305) Jackson | (306) Juneau | (307) Marathon | (308) Marquette |
| (309) Monroe | (310) Portage | (311) Waupaca | (312) Waushara |
| (313) Wood | | | |

Southwest Unit

| | | | |
|---|---|---|---|
| (401) Buffalo | (402) Crawford | (403) Dunn | (404) Grant |
| (405) Iowa | (406) La Crosse | (407) Lafayette | (408) Pepin |
| (409) Pierce | (410) Richland | (411) St. Croix | (412) Sauk |
| (413) Tempeleau | (414) Vernon | | |

Southeast Unit

| | | | |
|---|---|---|---|
| (501) Brown | (502) Calumet | (503) Columbia | (504) Dane |
| (505) Dodge | (506) Door | (507) Fond du Lac | (508) Green |
| (509) Green Lake | (510) Jefferson | (511) Kenosha | (512) Kewaunee |
| (513) Manitowoc | (514) Milwaukee | (515) Outagamie | (516) Ozaukee |
| (517) Racine | (518) Rock | (519) Sheboygen | (520) Walworth |
| (521) Washington | (522) Waukesha | (523) Winnebago | |

State = Minnesota State code = 27

Data base number and name (V) MINNOR

Aspen-Birch Unit

| | | | |
|---|---|---|---|
| (101) Carlton | (102) Cook | (103) Koochiching | (104) Lake |
| (105) St. Louis | | | |

Northern Pine Unit

| | | | |
|---|---|---|---|
| (201) Aitkin | (202) Becker | (203) Beltrami | (204) Cass |
| (205) Clearwater | (206) Crow Wing | (207) Hubbard | (208) Itasca |
| (209) Lake of the Woods | | (210) Mahnomen | (211) Roseau |
| (212) Wadena | | | |

Data base number and name (VI) MINSOU

Central Hardwood Unit

| | | | |
|---|---|---|---|
| (301) Anoka | (302) Benton | (303) Carver | (304) Chisago |
| (305) Dakota | (306) Douglas | (307) Fillmore | (308) Goodhue |
| (309) Hennepin | (310) Houston | (311) Isanti | (312) Kanabec |
| (313) Le Sueur | (314) Mille Lacs | (315) Morrison | (316) Olmsted |
| (317) Otter tail | (318) Pine | (319) Ramsey | (320) Rice |
| (321) Scott | (322) Sherburne | (323) Stearns | (324) Todd |
| (325) Wabasha | (326) Washington | (327) Winona | (328) Wright |

Prairie Unit

| | | | |
|---|---|---|---|
| (401) Big Stone | (402) Blue Earth | (403) Brown | (404) Chippewa |
| (405) Clay | (406) Cottonwood | (407) Dodge | (408) Fairbault |
| (409) Freeborn | (410) Grant | (411) Jackson | (412) Kandiyohi |
| (413) Kittson | (414) Lac qui Parle | (415) Lincoln | (416) Lyon |
| (417) McLeod | (418) Marshall | (419) Martin | (420) Meeker |
| (421) Mower | (422) Murray | (423) Nicollet | (424) Nobles |
| (425) Norman | (426) Pennington | (427) Pipestone | (428) Polk |
| (429) Pope | (430) Red Lake | (431) Redwood | (432) Renville |
| (433) Rock | (434) Sibley | (435) Steele | (436) Stevens |
| (437) Swift | (438) Traverse | (439) Waseca | (440) Watonwan |
| (441) Wilkin | (442) Yellow Medicine | | |

| ode | County name | Code | County name | Code | County name | Code | County name |
|---|---|---|---|---|---|---|---|

State = North Dakota  State code = 38
Data base number and name (VII) NDAKOTA

| | | | |
|---|---|---|---|
| 101) Adams | (102) Barnes | (103) Benson | (104) Billings |
| 105) Bottineau | (106) Bowman | (107) Burke | (108) Burleigh |
| 109) Cass | (110) Cavalier | (111) Dickey | (112) Divide |
| 113) Dunn | (114) Eddy | (115) Emmons | (116) Foster |
| 117) Golden Valley | (118) Grand Forks | (119) Grant | (120) Griggs |
| 121) Hettinger | (122) Kidder | (123) LaMoure | (124) Logan |
| 125) McHenry | (126) McIntosh | (127) McKenzie | (128) McLean |
| 129) Mercer | (130) Morton | (131) Mountrail | (132) Nelson |
| 133) Oliver | (134) Pembina | (135) Pierce | (136) Ramsey |
| 137) Ransom | (138) Renville | (139) Richland | (140) Rolette |
| 141) Sargent | (142) Slope | (143) Sheridan | (144) Sioux |
| 145) Stark | (146) Steele | (147) Stutsman | (148) Towner |
| 149) Traill | (150) Walsh | (151) Ward | (152) Wells |
| 153) Williams | | | |

State = South Dakota  State code = 46
Data base number and name (VIII) SDAKOTA

Eastern Unit

| | | | |
|---|---|---|---|
| (101) Aurora | (102) Beadle | (103) Bennet | (104) Bon Homme |
| (105) Brookings | (106) Brown | (107) Brule | (108) Buffalo |
| (109) Campbell | (110) Charles | (111) Clark | (112) Clay |
| (113) Codington | (114) Corson | (115) Custer | (116) Davison |
| (117) Day | (118) Deuel | (119) Dewey | (120) Douglas |
| (121) Edmunds | (122) Faulk | (123) Grant | (124) Gregory |
| (125) Haakon | (126) Hamlin | (127) Hand | (128) Hanson |
| (129) Hughes | (130) Hutchinson | (131) Hyde | (132) Jackson |
| (133) Jerauld | (134) Jones | (135) Kingsbury | (136) Lake |
| (137) Lincoln | (138) Lyman | (139) Marshall | (140) McCook |
| (141) McPherson | (142) Mead | (143) Mellette | (144) Miner |
| (145) Minnehaha | (146) Woody | (147) Pennington | (148) Perkins |
| (149) Potter | (150) Roberts | (151) Sanborn | (152) Shannon |
| (153) Spink | (154) Stanley | (155) Sully | (156) Todd |
| (157) Tripp | (158) Turner | (159) Union | (160) Walworth |
| (161) Washabaugh | (162) Yankton | (163) Ziebach | |

| Code | County name | Code | County name | Code | County name | Code | County name |
|---|---|---|---|---|---|---|---|

State = Nebraska State code = 31
Data base number and name (IX) NEBRASK

Eastern Unit

| | | | |
|---|---|---|---|
| (101) Adams | (102) Boone | (103) Buffalo | (104) Burt |
| (105) Butler | (106) Cass | (107) Cedar | (108) Clay |
| (109) Colfax | (110) Cuming | (111) Custer | (112) Dakota |
| (113) Dawson | (114) Dixon | (115) Dodge | (116) Douglas |
| (117) Fillmore | (118) Franklin | (119) Frontier | (120) Furnas |
| (121) Gage | (122) Gosper | (123) Greeley | (124) Hall |
| (125) Hamilton | (126) Harlan | (127) Hitchcock | (128) Howard |
| (129) Jefferson | (130) Johnson | (131) Kearney | (132) Lancaster |
| (133) Madison | (134) Merrick | (135) Nance | (136) Nemaha |
| (137) Nuckolls | (138) Otoe | (139) Pawnee | (140) Phelps |
| (141) Pierce | (142) Platte | (143) Polk | (144) Red Willow |
| (145) Richardson | (146) Saline | (147) Sarpy | (148) Saunders |
| (149) Seward | (150) Sherman | (151) Stanton | (152) Thayer |
| (153) Thurston | (154) Valley | (155) Washington | (156) Wayne |
| (157) Webster | (158) York | | |

Western Unit

| | | | |
|---|---|---|---|
| (201) Antelope | (202) Arthur | (203) Banner | (204) Blaine |
| (205) Box Butte | (206) Boyd | (207) Brown | (208) Chase |
| (209) Cherry | (210) Cheyenne | (211) Dawes | (212) Deuel |
| (213) Dundy | (214) Garden | (215) Garfield | (216) Grant |
| (217) Hayes | (218) Holt | (219) Hooker | (220) Keith |
| (221) Keya Paha | (222) Kimball | (223) Knox | (224) Lincoln |
| (225) Logan | (226) Loup | (227) McPherson | (228) Morrill |
| (229) Perkins | (230) Rock | (231) Scotts Bluff | (232) Sheridan |
| (233) Sioux | (234) Thomas | (235) Wheeler | |

Data base number and name (XI) IOWA

Northeast Unit

| | | | |
|---|---|---|---|
| (101) Allamakee | (102) Benton | (103) Black Hawk | (104) Bremer |
| (105) Buchanan | (106) Butler | (107) Cedar | (108) Chicksaw |
| (109) Clayton | (110) Clinton | (111) Delaware | (112) Dubuque |
| (113) Fayette | (114) Floyd | (115) Grundy | (116) Howard |
| (117) Jackson | (118) Johnson | (119) Jones | (120) Linn |
| (121) Mitchell | (122) Scott | (123) Tama | (124) Winneshiek |

Southeast Unit

| | | | |
|---|---|---|---|
| (201) Appanoose | (202) Boone | (203) Clark | (204) Dallas |
| (205) Davis | (206) Decatur | (207) Des Moines | (208) Guthrie |
| (209) Hamilton | (210) Hardin | (211) Henry | (212) Iowa |
| (213) Jasper | (214) Jefferson | (215) Keokuk | (216) Lee |
| (217) Louisa | (218) Lucas | (219) Madison | (220) Mahaska |
| (221) Marion | (222) Marshall | (223) Monroe | (224) Muscatine |
| (225) Polk | (226) Poweshiek | (227) Story | (228) Van Buren |
| (229) Wapello | (230) Warren | (231) Washington | (232) Wayne |
| (233) Webster | | | |

Southwest Unit

| | | | |
|---|---|---|---|
| (301) Adair | (302) Adams | (303) Audubon | (304) Carroll |
| (305) Cass | (306) Crawford | (307) Fremont | (308) Greene |
| (309) Harrison | (310) Mills | (311) Monona | (312) Montgomery |
| (313) Page | (314) Pottawattami | (315) Ringgold | (316) Shelby |
| (317) Taylor | (318) Union | (319) Woodbury | |

Northwest Unit

| | | | |
|---|---|---|---|
| (401) Buena Vista | (402) Calhoun | (403) Cerro Sordo | (404) Cherokee |
| (405) Clay | (406) Dickinson | (407) Emmet | (408) Franklin |
| (409) Hancock | (410) Humboldt | (411) Ida | (412) Kossuth |
| (413) Lyon | (414) O'Brien | (415) Osceola | (416) Palo Alto |
| (417) Plymouth | (418) Pocahontas | (419) Sac | (420) Sioux |
| (421) Winnebago | (422) Worth | (423) Wright | |

| Code | County name | Code | County name | Code | County name | Code | County name |
|---|---|---|---|---|---|---|---|

State = Missouri  State code = 29
Data base number and name (XII) MOZARK

Eastern Ozarks Unit

| | | | |
|---|---|---|---|
| (101) Bollinger | (102) Butler | (103) Carter | (104) Crawford |
| (105) Dent | (106) Iron | (107) Madison | (108) Oregon |
| (109) Reynolds | (110) Ripley | (111) St.Francois | (112) Shannon |
| (113) Washington | (114) Wayne | | |

Southwest Ozarks Unit

| | | | |
|---|---|---|---|
| (201) Barry | (202) Christian | (203) Douglas | (204) Howell |
| (205) McDonald | (206) Newton | (207) Ozark | (208) Stone |
| (209) Taney | (210) Texas | (211) Webster | (212) Wright |

Northwest Ozarks Unit

| | | | |
|---|---|---|---|
| (301) Benton | (302) Camden | (303) Cedar | (304) Dallas |
| (305) Hickory | (306) Laclede | (307) Maries | (308) Miller |
| (309) Morgan | (310) Phelps | (311) Polk | (312) Pulaski |
| (313) St.Clair | | | |

Data base number and name (XIII) MORBNW

Prairie Unit

| | | | |
|---|---|---|---|
| (401) Adair | (402) Andrew | (403) Atchison | (404) Audrain |
| (405) Barton | (406) Bates | (407) Buchanan | (408) Caldwell |
| (409) Carroll | (410) Cass | (411) Chariton | (412) Clark |
| (413) Clay | (414) Clinton | (415) Cooper | (416) Dade |
| (417) Daviess | (418) De Kalb | (419) Gentry | (420) Greene |
| (421) Grundy | (422) Harrison | (423) Henry | (424) Holt |
| (425) Jackson | (426) Jasper | (427) Johnson | (428) Knox |
| (429) Lafayette | (430) Lawrence | (431) Lewis | (432) Lincoln |
| (433) Linn | (434) Livingston | (435) Macon | (436) Marion |
| (437) Mercer | (438) Monroe | (439) Nodaway | (440) Pettis |
| (441) Pike | (442) Platte | (443) Putnam | (444) Rallis |
| (445) Randolph | (446) Ray | (447) Saline | (448) Schuyler |
| (449) Scotland | (450) Shelby | (451) Sullivan | (452) Vernon |
| (453) Worth | | | |

Riverborder Unit

| | | | |
|---|---|---|---|
| (501) Boone | (502) Callaway | (503) Cape Girardeau | (504) Cole |
| (505) Dunkin | (506) Franklin | (507) Gasconade | (508) Howard |
| (509) Jefferson | (510) Mississippi | (511) Moniteau | (512) Montgomery |
| (513) New Madrid | (514) Osage | (515) Pemiscot | (516) Perry |
| (517) Saint Charles | (518) Saint Louis | (519) Sainte Genevie | |
| (520) Scott | (521) Stoddard | (522) Warren | |
| (523) City of Saint Louis | | | |

Southern Unit

| | | | |
|---|---|---|---|
| (101) Alexander | (102) Franklin | (103) Gallatin | (104) Hamilton |
| (105) Hardin | (106) Jackson | (107) Johnson | (108) Massac |
| (109) Perry | (110) Pope | (111) Pulaski | (112) Randolph |
| (113) Saline | (114) Union | (115) White | (116) Williamson |

Claypan Unit

| | | | |
|---|---|---|---|
| (201) Bond | (202) Calhoun | (203) Clark | (204) Clay |
| (205) Clinton | (206) Crawford | (207) Cumberland | (208) Edwards |
| (209) Effingham | (210) Fayette | (211) Greene | (212) Jasper |
| (213) Jefferson | (214) Jersey | (215) Lawrence | (216) Macoupin |
| (217) Madison | (218) Marion | (219) Monroe | (220) Montgomery |
| (221) Richland | (222) St. Clair | (223) Shelby | (224) Wabash |
| (225) Washington | (226) Wayne | | |

| Code | County name | Code | County name | Code | County name | Code | County name |
|---|---|---|---|---|---|---|---|
| Prairie Unit | | | | | | | |
| (301) | Adams | (302) | Boone | (303) | Brown | (304) | Bureau |
| (305) | Carroll | (306) | Cass | (307) | Champaign | (308) | Christian |
| (309) | Coles | (310) | Cook | (311) | De Kalb | (312) | Dewitt |
| (313) | Douglas | (314) | Dupage | (315) | Edgar | (316) | Ford |
| (317) | Fulton | (318) | Grundy | (319) | Hancock | (320) | Henderson |
| (321) | Henry | (322) | Iroquois | (323) | Jo Daviess | (324) | Kane |
| (325) | Kankakee | (326) | Kendall | (327) | Knox | (328) | Lake |
| (329) | La Salle | (330) | Lee | (331) | Livingston | (332) | Logan |
| (333) | Macon | (334) | Marshall | (335) | Mason | (336) | McDonough |
| (337) | McHenry | (338) | McLean | (339) | Menard | (340) | Mercer |
| (341) | Morgan | (342) | Moultrie | (343) | Ogle | (344) | Peoria |
| (345) | Piatt | (346) | Pike | (347) | Putnam | (348) | Rock Island |
| (349) | Sangamon | (350) | Schuyler | (351) | Scott | (352) | Stark |
| (353) | Stephenson | (354) | Tazewell | (355) | Vermillion | (356) | Warren |
| (357) | Whiteside | (358) | Will | (359) | Winnebago | (360) | Woodford |

State = Indiana  State code = 18
Data base number and name  (XV) INDIANA

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Lower Wabash Unit | | | | | | | |
| (101) | Clay | (102) | Daviess | (103) | Gibson | (104) | Greene |
| (105) | Knox | (106) | Martin | (107) | Parke | (108) | Pike |
| (109) | Posey | (110) | Putnam | (111) | Sullivan | (112) | Vanderburg |
| (113) | Vermillion | (114) | Vigo | | | | |
| Knobs Unit | | | | | | | |
| (201) | Brown | (202) | Clark | (203) | Crawford | (204) | Dubois |
| (205) | Floyd | (206) | Harrison | (207) | Jackson | (208) | Lawrence |
| (209) | Monroe | (210) | Morgan | (211) | Orange | (212) | Owen |
| (213) | Perry | (214) | Scott | (215) | Spencer | (216) | Warrick |
| (217) | Washington | | | | | | |
| Upland Flats Unit | | | | | | | |
| (301) | Dearborn | (302) | Fayette | (303) | Franklin | (304) | Jennings |
| (305) | Jefferson | (306) | Ohio | (307) | Ripley | (308) | Switzerland |
| (309) | Union | | | | | | |
| Northern Unit | | | | | | | |
| (401) | Adams | (402) | Allen | (403) | Bartholomew | (404) | Benton |
| (405) | Blackford | (406) | Boone | (407) | Carroll | (408) | Cass |
| (409) | Clinton | (410) | Decatur | (411) | De Kalb | (412) | Delaware |
| (413) | Elkhart | (414) | Fountain | (415) | Fulton | (416) | Grant |
| (417) | Hamilton | (418) | Hancock | (419) | Hendricks | (420) | Henry |
| (421) | Howard | (422) | Huntington | (423) | Jasper | (424) | Jay |
| (425) | Johnson | (426) | Kosciusko | (427) | La Grange | (428) | Lake |
| (429) | La Porte | (430) | Madison | (431) | Marion | (432) | Marshall |
| (433) | Miami | (434) | Montgomery | (435) | Newton | (436) | Noble |
| (437) | Porter | (438) | Pulaski | (439) | Randolph | (440) | Rush |
| (441) | St. Joseph | (442) | Shelby | (443) | Starke | (444) | Steuben |
| (445) | Tippecanoe | (446) | Tipton | (447) | Wabash | (448) | Warren |
| (449) | Wayne | (450) | Wells | (451) | White | (452) | Whitley |

Codes for variables R4 PICODE, R5 PISTRAT

The use of these variables was described in section I sampling and estimation procedures. The actual codes vary from State to State. PICODE generally indicates a broad land use category such as (10 or 20) forest, (50 or 60) nonforest, (30) questionable, and (80 or 90) water. PISTRAT indicates a broad forest type and stand-size density class using codes similar to but not identical to variables C4 FORTCUR and R21 SZDNCUR.

Codes for variable R9 UTREND

The first two digits describe the land class change that took place between surveys or the change since the date of photography. Codes to be used for the first two digits are divided into the following two categories:

When land class is not commercial forest on both occasions:

First two digits--The first digit is the code for the present land class. The second digit is the code of the land class at the time of the last survey or the date of photography.

| Code | Land Use Class |
|---|---|
| 1 | Commercial forest |
| 2 | Productive reserved forest |
| 3 | Unproductive forest |
| 4 | Cropland |
| 5 | Pasture, rangeland |
| 6 | Idle farmland |
| 7 | Wooded pasture |
| 8 | Urban, recreation, wooded strips, rights of way, other |
| 9 | Water and marsh |

For commercial forest land on both occasions:

First two digits--The codes are used to indicate any major changes in the stand since the last survey or the date of photography.

If there has been a disturbance in the sampling area since the date of photography but the forest type or stand-size class has not changed, it will be recorded as no change. This disturbance could be recorded under stand history.

| Code | Stand changes |
|---|---|
| 10 | No change |
| 01 | Forest type change |
| 02 | Stand-size change |
| 03 | Forest type and size change |

The third digit indicates the process that caused the change and uses one of the following codes:

| Code | Cause of Land Use Change |
|---|---|
| 0 | No change |
| 1 | Definition |
| 2 | Legislation |
| 3 | Natural |
| 4 | Herbicide |
| 5 | Clearing (land cleared by mechanical or hand means but timber not utilized) |
| 6 | Clearcut (includes land clearing where timber is utilized) |

7 Partial cut
8 Planting
9 Other man (includes fencing to exclude livestock)

Examples: 1. Commercial forest land (Red Pine Plantation) now, on the photo was idle farmland, use trend code 168.

2. Marsh without trees now, on photo was marsh with trees, use trend code 990.

3. Cleared powerline right of way now, on photo was commercial forest land. Timber utilized, use trend code 816.

Codes for variables R10 STDHIST, R11 PRJSTDH

Stand history reflects the kind of disturbance on five or more of the sample points within the last 20 years.

Explain the kind and extent of any disturbance in the "Notes" on the back of the plot sheet. Use the following 2-digit code to record stand history:

First digit (what happened)

1 No disturbance
2 Timber stand improvement
3 Clearcut
4 Partial harvest cut
5 Natural--fire, insects, disease
6 Man caused--drainage, spraying
7 Planting of forest land
8 Planting of nonforest land
9 Natural regeneration of non-forest land

Second digit (how long ago)

0 No disturbance
1 1-4 years
2 5-10 years
3 11-15 years
4 15-20 years

Codes for variable R35 AREACOND

(10) Areas fully stocked with desirable trees but not overstocked.

(20) Areas fully stocked with desirable trees but overstocked with all live trees.

(30) Areas medium to fully stocked with desirable trees and with less than 30 percent of the area controlled by other trees and/or inhibiting vegetation or surface conditions that will prevent occupancy by desirable trees.

(40) Areas medium to fully stocked with desirable trees and with 30 percent or more of the area controlled by other trees and/or conditions that ordinarily prevent occupancy by desirable trees.

(50) Areas poorly stocked with desirable trees but fully stocked with growing-stock trees.

(60) Areas poorly stocked with desirable trees but with medium to full stocking of growing-stock trees.

(70) Areas poorly stocked with desirable trees and poorly stocked with growing-stock trees.

Codes for variable R10 CAVIT

At each sample point, each live tree 5.0 inches DBH and larger is examined for cavities used for nesting, resting or storage by birds or mammals. For the largest cavity a 2-digit code is recorded to indicate the size of the cavity entrance hole and location of the cavity in the tree. The first digit indicates the cavity hole size, the second digit indicates the location of the cavity.

To qualify as a cavity, the entrance hole must be 1.0 inch or larger in the main stem, fork, or larger limb. (A large limb must be greater than 8.0 inch in diameter o.b.). If no cavity is present, this item is zero or missing.

missing.

| First digit | | Second digit | |
|---|---|---|---|
| Code | Size of opening (inches) | Code | Location of cavity (feet) |
| 1 | 1 | 1 | 0-1 |
| 2 | 2 | 2 | 2-5 |
| 3 | 3 | 3 | 6-9 |
| 4 | 4 | 4 | 10-19 |
| 5 | 5 | 5 | 20-29 |
| 6 | 6 | 6 | 30-39 |
| 7 | 7 | 7 | 40-49 |
| 8 | 8 | 8 | 50-59 |
| 9 | 9+ | 9 | 60+ |

Codes for variables C4 FORTCUR, R20 FORTPRJ, R23 FORTORG

(For detailed definitions of these forest types see page 14 )

States = Michigan, Wisconsin, Minnesota
Data base numbers and names (I)MICHUP, (II)MICHLP, (III)WISNOR, (IV)WISSOU, (V)MINNOR, (VI)MINSOU

| Code | Forest type | Code | Forest type |
|---|---|---|---|
| ( 1) | Jack pine | ( 2) | Red pine |
| ( 3) | White pine | ( 6) | Exotic |
| (12) | Black spruce | (13) | Balsam fir |
| (14) | Northern white-cedar | (15) | Tamarack |
| (16) | White spruce | (50) | Oak-hickory |
| (70) | Elm-ash-soft maple | (80) | Maple-birch |
| (91) | Aspen | (92) | Paper birch |

States = North Dakota, South Dakota, Nebraska, Kansas
Data base numbers and names (VII)NDAKOTA, (VIII)SDAKOTA, (IX)NEBRASK, (X)KANSAS

| | | | |
|---|---|---|---|
| (42) | Cedar-hardwood | (50) | Oak-hickory |
| (51) | Post-blackjack oak | (59) | Upland plains hwds. |
| (70) | Elm-ash-cottonwood | (74) | Willow |
| (73) | Cottonwood | (87) | Upland elm-ash-loc. |
| (77) | Lowland plains hwds. | | |

State = Iowa
Data base number and name (XI)IOWA

| | | | |
|---|---|---|---|
| (42) | E.redcedar-hardwood | (50) | Oak-hickory |
| (54) | White oak | (59) | Bur oak |
| (70) | Elm-ash-cottonwood | (73) | Cottonwood |
| (80) | Maple-basswood | (91) | Aspen |

State = Missouri
Data base numbers and names (XII)MOZARK, (XIII)MORBNW

| Code | Forest type | Code | Forest type |
|---|---|---|---|
| (32) | Shortleaf pine | (35) | Eastern redcedar |
| (42) | Redcedar-hardwood | (44) | Shortleaf pine-oak |
| (51) | Post-blackjack oak | (54) | White oak |
| (53) | Black-scarlet oak | (70) | Elm-ash-cottonwood |
| (60) | Oak-gum-cypress | | |
| (73) | Cottonwood | | |
| (80) | Maple-beech | | |

States = Illinois,Indiana
Data base numbers and names (XIV)ILLINI, (XV)INDIANA

| | | | |
|---|---|---|---|
| (30) | Loblolly-shortleaf pine | (40) | Oak-pine |
| (50) | Oak-hickory | (60) | Oak-gum-cypress |
| (70) | Elm-ash-cottonwood | | |
| (80) | Maple-beech-birch | | |
| (90) | Aspen-birch | | |

Codes for species variables R16 CUNSPP, R28 SISPP, R4 SPECIES, R3 SPPSI

| Code | Commercial species Common name | Scientific name |
|---|---|---|
| 012 | Balsam fir | Abies balsamea var. |
| 068 | Eastern redcedar | Juniperus virginiana |
| 070 | European larch | Larix decidua |
| 071 | Tamarack | Larix laricina |
| 091 | Norway spruce | Picea abies |
| 093 | Engelmann spruce | Picea engelmannii |
| 094 | White spruce | Picea glauca |
| 095 | Black spruce | Picea mariana |
| 096 | Col. blue spruce | Picea pungens |
| 105 | Jack pine | Pinus banksiana |
| 113 | Limber pine | Pinus flexilis |
| 122 | Ponderosa pine | Pinus ponderosa |
| 125 | Red pine | Pinus resinosa |
| 129 | White pine | Pinus strobus |
| 130 | Scotch pine | Pinus sylvestris |
| 133 | Austrian pine | Pinus nigra |
| 202 | Douglas fir | Pseudotsuga menziesii |
| 241 | N. white-cedar | Thuja occidentalis |
| 261 | Hemlock | Tsuga canadensis |
| 300 | Acacia | Acacia spp. |
| 313 | Boxelder | Acer negundo |
| 314 | Black maple | Acer nigrum |
| 316 | Red maple | Acer rubrum |
| 317 | Silver maple | Acer saccharinum |
| 318 | Sugar maple | Acer saccharum |
| 331 | Ohio buckeye | Aesculus glabra |
| 371 | Yellow birch | Betula alleghaniensis |
| 372 | Sweet birch | Betula lenta |
| 373 | River birch | Betula nigra |
| 374 | Water birch | Betula occidentalis |
| 375 | Paper birch | Betula papyrifera |
| 402 | Bitternut hickory | Carya cordiformis |
| 403 | Pignut hickory | Carya glabra |
| 404 | Pecan | Catya illinoensis |
| 405 | Shellbark hickory | Carya laciniosa |
| 407 | Shagbark hickory | Carya ovata |
| 408 | Black hickory | Carya texana |
| 409 | Mockernut hickory | Carya tomentosa |
| 421 | American chestnut | Castanea dentata |
| 452 | Northern catalpa | Catalpa speciosa |
| 461 | Sugarberry | Celtis laevigata |
| 462 | Hackberry | Celtis occidentalis |
| 491 | Flowering dogwood | Cornus florida |
| 521 | Persimmon | Diospyros virginiana |
| 531 | Beech | Fagus grandifolia |
| 541 | White ash | Fraxinus americana |
| 543 | Black ash | Fraxinus nigra |
| 544 | Green ash | Fraxinus pennsylvanica |
| 546 | Blue ash | Fraxinus quadrangulata |
| 552 | Honeylocust | Gleditsia triacanthus |
| 571 | Ky. coffee tree | Gymnocladus dioicus |
| 601 | Butternut | Juglans cinerea |
| 602 | Black walnut | Juglans nigra |
| 621 | Yellow poplar | Liriodendron tulipifera |
| 660 | Apple | Malus spp. |
| 681 | White mulberry | Morus alba |
| 682 | Red mulberry | Morus rubra |
| 693 | Black tupelo | Nyssa sylvatica |
| 731 | Sycamore | Platanus occidentalis |
| 741 | Balsam poplar | Populus balsamifera |
| 742 | East. cottonwood | Populus deltoides |
| 743 | Bigtooth aspen | Populus grandidentata |
| 745 | Plains cottonwood | Populus sargentii |
| 746 | Quaking aspen | Populus tremuloides |
| 752 | Silver poplar | Populus alba |

| Code | Common name | Scientific name |
|---|---|---|
| 753 | Narrowleaf ctnwd. | Populus angustifolia |
| 762 | Black cherry | Prunus serotina |
| 802 | White oak | Quercus alba |
| 804 | Swamp white oak | Quercus bicolor |
| 806 | Scarlet oak | Quercus coccinea |
| 809 | Northern pin oak | Quercus ellipsoidalis |
| 817 | Shingle oak | Quercus imbricaria |
| 823 | Bur oak | Quercus macrocarpa |
| 824 | Blackjack oak | Quercus marilandica |
| 826 | Chinkapin oak | Quercus muehlenbergii |
| 830 | Pin oak | Quercus palustris |
| 832 | Chestnut oak | Quercus prinus |
| 833 | Northern red oak | Quercus rubra |
| 834 | Shumard oak | Quercus shumardii |
| 835 | Post oak | Quercus stellata |
| 837 | Black oak | Quercus velutina |
| 901 | Black locust | Robinia pseudoacacia |
| 922 | Black willow | Salix nigra |
| 931 | Sassafras | Sassafras albidum |
| 951 | American basswood | Tilia americana |
| 972 | American elm | Ulmus americana |
| 974 | Siberian elm | Ulmus pumila |
| 975 | Slippery elm | Ulmus rubra |
| 977 | Rock elm | Ulmus thomasii |

Noncommercial species

| | | |
|---|---|---|
| 066 | Rocky mt. juniper | Juniperus scorplorum |
| 315 | Striped maple | Acer pensylvanicum |
| 319 | Mountain maple | Acer spicatum |
| 321 | Rocky mt. maple | Acer glabrum |
| 333 | Western buckeye | Aesculus glabra |
| 341 | Ailanthus | Ailanthus altissima |
| 391 | Am. hornbeam | Carpinus caroliniana |
| 471 | Eastern redbud | Cercis canadensis |
| 500 | Hawthorn | Crataegus species |
| 641 | Osage orange | Maclura pomifera |
| 701 | Ironwood | Ostrya virginiana |
| 761 | Pincherry | Prunus pensylvanica |
| 763 | Chokecherry | Prunus virginiana |
| 765 | Canada plum | Prunus nigra |
| 766 | Wild plum | Prunus americana |
| 851 | Mountain ash | Sorbus |
| 921 | Peachleaf willow | Salix amygdaloides |
| 923 | Diamond willow | Salix eriocephala |

Codes for record type 4 variable R2 SHRUBSP tall woody shrubs

| Code | Common name | Scientific name |
|---|---|---|
| 351 | Red alder | Alnus rubra |
| 352 | Green alder | Alnus crispa |
| 353 | Speckled alder | Alnus rugosa |
| 380 | Swamp birch | Betula pumila |
| 381 | Dwarf birch | Betula glandulosa |
| 463 | Dwarf hackberry | Celtis tenuiflolia |
| 490 | Dogwood | Cornaceae |
| 592 | Black alder | Ilex verticillata |
| 593 | Mountain holly | Nemopanthus mucronata |
| 594 | Deciduous holly | Ilex decidua |
| 603 | Spice bush | Lindera benzoin |
| 764 | Sandcherry | Prunus pumila |
| 767 | Pawpaw | Asimina triloba |
| 768 | Devils wlk. stick | Asimina triloba |
| 769 | Chickasaw plum | Prunus angustifolia |
| 850 | Sweet gale | Myrica gale |
| 852 | Common barberry | Berberis vulgaris |

| | | |
|---|---|---|
| 853 | Witch hazel | Hamamelis virginiana |
| 854 | Common ninebark | Physocarpus opulifolius |
| 855 | Juneberry | Amelanchier |
| 856 | Beaked hazenut | Corylus cornuta |
| 857 | Prickly ash | Zanthoxylum americanum |
| 859 | Buckthorn | Rhamnus |
| 860 | Mountain laurel | Ceanothus sanguineus |
| 861 | Leatherwood | Dirca palustris |
| 862 | Vibernum | Viburnum |
| 863 | Elderberry | Sambucus |
| 864 | Sumac | Rhus |
| 869 | Shrubby trefoil | Ptelea trifoliata |
| 880 | Buffaloberry | Shepherdia canadensus |
| 903 | Poison sumac | Rhus vernix |
| 907 | New Jersey tea | Ceanothus americanus |
| 912 | Buttonbush | Cephalanthus occidentalis |
| 913 | Russian olive | Elaeagnus |
| 915 | Wild crabapple | Pyrus ioensis |
| 916 | Lead plant | Amorpha |
| 917 | Wahoo | Euonymus atropurpurea |
| 919 | Soapberry | Sapindus drummondii |
| 920 | Willow | Salicaceae |
| 925 | Tamarisk | Amarix gallica |
| 926 | Buckthorn | Umelia lanuginosa |
| 927 | Rabbitbush | Chrysothamnus pulchell |
| 997 | Other species | (tall, perennials) |

Codes for record type 5 variable R2 SHUSPL low herbaceous shrubs.

| | | |
|---|---|---|
| 059 | Creeping juniper | Juniperus horizontalis |
| 069 | Common juniper | Juniperus communis |
| 230 | Yew | Taxus canadensis |
| 590 | Holly | Ilex |
| 712 | Virginia creeper | Parthenocissus |
| 748 | Spirea | Spiraea |
| 749 | Labrador tea | Ledum groenlandicum |
| 750 | Leatherleaf | Chamaedaphne calyculat |
| 751 | Bog laurel | Kalmia polifolia |
| 849 | Sweetfern | Comptonia peregrina |
| 865 | Gooseberry | Ribes |
| 866 | Chokeberry | Aronia |
| 867 | Rasp.-blackberry | Rubus |
| 868 | Rose | Rosa |
| 870 | Am. bladdernut | Staphylea trifolia |
| 871 | Willow herb | Decodon verticillatus |
| 872 | Privet andromeda | Lyonia liqustrina |
| 873 | Black huckleberry | Gaylussacia baccata |
| 874 | Bil.-blueberry | Vaccinium |
| 875 | Bush honeysuckle | Diervilla lonicera |
| 876 | Honeysuckle | Lonicera |
| 877 | Buckbrush | Symphoricarpus |
| 878 | Shrb. cinquefoil | Potentilla fruticosa |
| 902 | Poison ivy | Rhus radicans |
| 908 | St.johns wort | Hypericum |
| 909 | Bearberry | Arctosttaphylos uva-ur |
| 918 | Bittersweet | Celastrus scandens |
| 924 | Raccoon grape | Ampelopsis cordata |
| 929 | Sandhill sage | Artemisia filifolia |
| 932 | Greenbriar | Smilax |
| 978 | Bog rosemary | Andromeda glaucophylla |
| 982 | Grape | Vitis |
| 983 | Clematis | Vitis |
| 984 | Strawberry | Fragaria spp. |
| 998 | Other species | (other perennials) |

## APPENDIX B.--GLOSSARY OF TERMS

Acceptable trees.--Growing-stock trees of commercial species that meet specified standards of size and quality but do not qualify as desirable trees.

Basal area.--The area in square feet of the cross section at breast height of a single tree. When the basal area of all trees in a stand are summed, the result is usually expressed as square feet of basal area per acre.

Biomass.--The above-ground volume of all live trees (including bark and foliage) reported in green tons. Biomass is made up of five components:

Growing-stock bole.--Biomass of a growing-stock tree from a 1-foot stump to a variable 4-inch top.

Growing-stock tops and limbs.--Biomass of a growing-stock tree from a 1-foot stump minus the growing-stock bole.

Cull bole.--Biomass of a cull tree from a 1-foot stump to a variable 4-inch top.

Cull tops and limbs.--Biomass of a cull tree from a 1-inch stump minus the cull bole.

1- to 5-inch trees.--Biomass of all live trees 1 to 5 inches in diameter at breast height.

Commercial forest land.--Forest land producing or capable of producing crops of industrial wood and not withdrawn from timber utilization. (Note: Areas qualifying as commerial forest land are capable of producing more than 20 cubic feet per acre per year of annual growth under management. Currently inaccessible and inoperable areas are included, except when the areas involved are small and unlikely to become suitable for producing industrial wood in the foreseeable future.) Also see definition of pastured commercial forest land.

Commercial species.--Tree species presently or prospectively suitable for industrial wood products. (Note: Excludes species of typically small size, poor form, or inferior quality such as hophornbeam and hawthorn.)

County and municipal land.--Land owned by counties and local public agencies or municipalities, or land leased to these governmental units for 50 years or more.

Cull.--Portions of a tree that are unusable for industrial wood products, because of rot, form, or other defect.

Desirable tree.--Growing-stock tree that has no serious defects in quality limiting present or prospective use, having relatively high vigor, and containing no pathogens that may kill or seriously deteriorate it before rotation age. These trees would be favored by forest managers in silvicultural operations.

Diameter classes.--A classification of trees based on diameter outside bark, measured at breast height (4.5 feet above the ground). (Note: d.b.h. is the common abbreviation for diameter at breast height. Two-inch diameter classes are commonly used in Forest Survey, with the even inch the approximate midpoint for a class. For example, the 6-inch class includes trees 5.0 through 6.9 inches d.b.h.)

Farm.--Either a place operated as a unit or 10 or more acres from which the sale of agricultural products totals $50 or more annually, or a place operated as a unit of less than 10 acres from which the sale of agricultural products for a year amounts to at least $250. Places having less than the $50 or $250 minimum estimated sales in a given year are also counted as farms if they can normally be expected to produce goods in sufficient quantity to meet the requirements of the definition.

Farmer-owned land.--Land owned by farm operators. (Note: Excludes land leased by farm operators from nonfarm owners, such as railroad companies and States.)

Forest land.--Land at least 16.7 percent stocked by forest trees of any size, or formerly having had such tree cover, and not currently developed for nonforest use. (Note: Stocking is measured by comparison of basal area and/or number of trees, by age or size and spacing with specified standards.) The minimum area for classification of forest land is one acre. Roadside, streamside, and shelterbelt strips of timber must have a a crown width of at least 120 feet to qualify as forest land. Unimproved roads and trails, streams, or other bodies of water or clearings in forest areas shall be classed as forest if less than 120 feet wide. Also see definitions for land area, commercial forest land, noncommercial forest land, productive-reserved forest land, stocking, unproductive forest land, and water.

Forest industry land.--Land owned by companies or individuals operating primary wood-using plants.

Forest trees.--Woody plants having a well-developed stem and usually more than 12 feet tall at maturity.

Forest type.--A classification of forest land based on the species forming a plurality of live tree stocking. Major forest types are:

Jack pine.--Forests in which jack pine comprises a plurality of the stocking. (Common associates include eastern white pine, red pine, aspen, birch, and maple.)

Red pine.--Forests in which red pine comprises a plurality of the stocking. (Common associates include eastern white pine, jack pine, aspen, birch, and maple.)

White pine.--Forests in which eastern white pine comprises a plurality of the stocking. (Common associates include red pine, jack pine, aspen, birch, and maple.)

Balsam fir.--Forests in which balsam fir and white spruce comprise a plurality of stocking with balsam fir the most common. (Common associates include white spruce, aspen, maple, birch, northern white-cedar, and tamarack.)

White spruce.--Forests in which white spruce and balsam fir comprise a plurality of the stocking with white spruce the most common. (Common associates include balsam fir, aspen, maple, birch, northern white-cedar, and tamarack.)

Black spruce.--Forests in which swamp conifers comprise a plurality of the stocking with black spruce the most common. (Common associates include tamarack and northern white-cedar.)

Northern white-cedar.--Forests in which swamp conifers comprise a plurality of the stocking with northern white-cedar the most common. (Common associates include tamarack and black spruce.)

Tamarack.--Forests in which swamp conifers comprise a plurality of the stocking with tamarack the most common. (Common associates include black spruce and northern white-cedar.)

Shortleaf pine.--Forests in which shortleaf pine comprises a plurality of the stocking. (Common associates include oak, hickory, and gum.)

Loblolly-shortleaf pine.--Forests in which loblolly, shortleaf, and Virginia pines, singly or in combination, comprise a plurality of the stocking. (Common associates include gum, hickory, sassafras, and yellow-poplar.)

Eastern redcedar.--Forests in which eastern redcedar comprises a plurality of the stocking. (Common associates are oak and hickory.)

Eastern redcedar-hardwood.--Forests in which hardwoods comprise a plurality of the stocking but in which eastern redcedar comprises 25 percent or more of the stocking. Found on dry uplands, usually abandoned pastures or fields.

Shortleaf pine-oak.--Forests in which upland oaks comprise a plurality of the stocking, but in which shortleaf pine comprises 25 to 50 percent of the stocking.

Oak-hickory.--Forests in which northern red oak, white oak, bur oak, or hickories, singly or in combination, comprise a plurality of the stocking. (Common associates include jack pine, beech, yellow-poplar, elm, and maple.)

Post-blackjack oak.--Forests in which post oak or blackjack oak, singly or in combination, comprise a majority of the stocking. Occurs on dry uplands and ridges.

Black-scarlet oak.--Forests in which upland oaks or hickory, singly or in combination, comprises a plurality of the stocking except where shortleaf pine or redcedar comprises 25 to 50 percent, or where white oak or post and blackjack oak comprise a plurality. (Common associates include yellow-poplar, elm, maple, and black walnut.)

White oak.--Forests in which white oak comprises more than 50 percent of the primary typing species for the oak-hickory type. (Common associates are black oak, northern red oak, bur oak, shagbark and bitternut hickories, white ash, and bigtooth aspen.)

Bur oak.--Forests in which bur oak comprises more than 50 percent of the stocking of the primary typing species for the oak-hickory type. (Common associates are northern pin oak, northern red oak, white oak, black oak, basswood, American elm, green ash, boxelder, hackberry, cottonwood, and hophornbeam.)

Upland plains hardwoods.--Forests in which black walnut, hackberry, and bur oak, singly or in combination, comprise a plurality of the stocking. Commonly found on slopes and uplands.

Oak-gum-cypress.--Bottomland forests in which bottomland oaks such as pin, swamp white, and shingle oaks, along with tupelo, blackgum, sweetgum, and cypress, singly or in combination, comprise a plurality of the stocking. (Common associates include cottonwood, willow, ash, elm, hackberry, and maple.)

Elm-ash-soft maple.--Forests in which lowland elm, ash, cottonwood, and red maple, singly or in combination, comprise a plurality of the stocking. (Common associates include birches, spruce, and balsam fir.)

Elm-ash-cottonwood.--Lowland forests in which elm, ash, cottonwood, and willow, singly or in combination, comprise a plurality of the stocking, except for those in which cottonwood or willow comprise a majority of the stocking. Found on first or second bottoms of major streams.

Cottonwood.--Forests in which cottonwood comprises a majority of the stocking.

Willow.--Forests in which willow comprises a majority of the stocking.

Lowland plains hardwoods.--Forests in which black walnut, hackberry, bur oak, soft maple, and boxelder, singly or in combination comprise a plurality of the stocking. Commonly found in coves and bottomlands.

Maple-birch.--Forests in which sugar maple, basswood, yellow birch, upland American elm, and red maple, singly or in combination, comprise a plurality of the stocking. (Common associates include white pine, elm, hemlock, and basswood.)

Maple-beech.--Forests in which hard maple or beech, singly or in combination, comprises a plurality of the stocking. (Common associates include elm and basswood.)

Hard maple-basswood.--Forests in which sugar maple or basswood, singly or in combination, comprise a plurality of the stocking. (Common associates are American elm, green ash, yellow birch, white pine, and northern red oak.)

Upland elm-ash-locust.--Upland forests in which elm, ash, and honeylocust, singly or in combination, comprise a plurality of the stocking. Includes shelterbelts and windbreaks on sites drier than those commonly associated with lowland species.

Aspen.--Forests in which quaking aspen or bigtooth aspen, singly or in combination, comprise a plurality of the stocking. (Common associates include balsam poplar, balsam fir, and paper birch.) and red maple, singly

or in combination, comprise a plurality of the stocking. (Common associates include birches, spruce, and balsam fir.)

Paper birch.--Forests in which paper birch comprises a plurality of the stocking. (Common associates include maple, aspen, and balsam fir.)

Exotic.--Forests in which species not native to the State comprise a plurality of the stocking. (Mostly Scotch pine plantations.)

Gross area.--The entire area of land and water as determined by the Bureau of the Census, 1970.

Growing-stock trees.--Live trees of commercial species qualifying as desirable and acceptable trees. (Note: Excludes rough, rotten, and dead trees.)

Growing-stock volume.--Net volume in cubic feet of growing-stock trees 5.0 inches d.b.h. and over, from a 1-foot stump to a minimum 4.0 inch top diameter outside bark of the central stem or to the point where the central stem breaks into limbs. Cubic feet can be converted to cords by dividing by 79 cubic feet per solid wood cord.

Hardwoods.--Dicotyledonous trees, usually broad-leaved and deciduous.

Idle farmland.--Includes former cropland, orchards, improved pastures, and farm sites not tended within the past 2 years and presently less than 16.7 percent stocked with trees.

Improved pasture.--Land currently improved for grazing by cultivating, seeding, irrigating or clearing of trees or brush and less than 16.7 percent stocked with live trees.

Indian land.--All lands held in trust by the United States for individual Indians or tribes, or all lands, titles to which are held by individual Indians or tribes, subject to Federal restrictions against alienation.

Land area.--A. Bureau of the Census. The area area of dry land and land temporarily or partly covered by water such as marshes, swamps, and river flood plains (omitting tidal flats below mean high tide); streams, sloughs, estuaries, and canals less than one-eighth of a statute mile wide; and lakes, reservoirs, and ponds less than 40 acres in area.

B. Forest Inventory and Analysis.--The same as the Bureau of the Census, except minimum width of streams, etc., is 120 feet and minimum size of lakes, etc., is 1 acre.

Live trees.--Growing-stock, rough, and rotten trees 1 inch d.b.h. and larger.

Log grades.--A classification of logs based on external characteristics as indicators of quality or value. (See Appendix for specific grading factors used.)

Logging residues.--The unused growing stock portions of trees cut or killed by logging.

Maintained road.--Any road, hard-topped or other surfaces, that is plowed or graded at least once a year. Includes rights-of-way that are cut or treated to limit herbaceous growth.

Marsh.--Nonforest land that characteristically supports low, generally herbaceous or shrubby vegetation and that is intermittently covered with water.

Merchantable.--Refers to a pulpwood or saw log section that meets pulpwood or saw log specfications, respectively.

Miscellaneous federal land.--Federal land other than National Forest, land administered by the Bureau of Land Management, and Indian land.

Miscellaneous private land.--Privately owned land other than forest-industry and farmer-owned land.

Mortality.--The volume of sound wood in growing-stock and sawtimber trees that die annually.

National Forest land.--Federal land that has been legally designated as National Forest or purchase units, and other land administered by the USDA Forest Service.

Net annual growth of growing-stock.--The annual change in volume of sound wood in live sawtimber and poletimber trees and the total volume of trees entering these classes through ingrowth, less volume losses resulting from natural causes.

Net annual growth of sawtimber.--The annual change in the volume of live sawtimber trees and the total volume of trees reaching sawtimber size, less volume losses resulting from natural causes.

Net volume.--Gross volume less deductions for rot, sweep, or other defect affecting use for timber products.

Noncommercial forest land.--(a) Unproductive forest land and (b) productive-reserved forest land.

Noncommercial species.--Tree species of typically small size, poor form, or inferior quality that normally do not develop into trees suitable for industrial wood products.

Nonforest land.--Land that has never supported forests, and land formerly forested where use for timber management is precluded by development for other uses. (Note: Includes areas used for crops, improved pasture, residential areas, city parks, improved roads of any width and adjoining clearings, powerline clearings of any width, and 1- to 40-acre areas of water classified by the Bureau of the Census as land. If intermingled in forest areas, unimproved roads and nonforest strips must be more than 120 feet wide and more than 1 acre in area to qualify as nonforest land.)

a. Nonforest land without trees.--Nonforest land with no live trees present.

b. Nonforest land with trees.--Nonforest land with one or more trees per acre at least 5 inches d.b.h.

Nonstocked land.--Commercial forest land less than 16.7 percent stocked with growing-stock trees.

Other removals.--Growing-stock trees removed but not utilized for products, or trees left standing but "removed" from the commercial forest land classification by land use change. Examples are removals from cultura operations such as timber stand improvement work, land clearing, and changes in land use.

Ownership.--Property owned by one owner, regardless of the number of parcels in a specified area.

Ownership size class.--The amount of commmercial forest land owned by one owner, regardless of the number of parcels.

Owner tenure.--The length of time a property has been held by the owner.

Physiographic class.--A measure of soil and water conditions that affect tree growth on a site. The physiographic classes are:

Xeric sites.--Very dry soils where excessive drainage seriously limits both growth and species occurrence. Example: sandy jack pine plains.

Xeromesic sites.--Moderately dry soils where excessive drainage limits growth and species occurrence to some extent. Example: dry oak ridge.

Mesic sites.--Deep, well-drained soils. Growth and species occurrence are limited only by climate.

Hydromesic sites.--Moderately wet soils where insufficient drainage or infrequent flooding limits growth and species occurrence to some extent. Example: better drained bottomland hardwood sites.

Hydric sites.--Very wet sites where excess water seriously limits both growth and species occurrence. Example: extra spruce wet, frequently flooded river bottoms and spruce bogs.

Plant byproducts.--Plant residues used for products such as mulch, pulp chips, and fuelwood.

Plant residues.--Wood and bark materials generated at manufacturing plants during production of other products.

Poletimber stands.--(See stand-size class.)

Poletimber trees.--Growing-stock trees of commercial species at least 5 inches d.b.h. but smaller than sawtimber

Productive-reserved forest land.--Forest land sufficiently productive to qualify as commercial forest land but withdrawn from timber utilization through statute, administration regulation, designation, or exclusive use for Christmas tree production, as indicated by annual shearing.

Productive-deferred.--Forest land sufficiently productive to qualify as commercial forest land but presently withdrawn from timber utilization because it is being considered for possible inclusion into the Wilderness system.

Rotten trees.--Live trees of commercial species that do not contain at least one 12-foot saw log or two saw logs 8 feet or longer, now or prospectively, and/or do not meet regional specifications cations for freedom from defect primarily because of rot; that is, when more than 50 percent of extra cull volume in a tree is rotten.

Rough trees.--(a) Live trees of commercial species that do not contain at least one merchantable 12-foot saw log or two saw logs 8 feet or longer, now or prospectively, and/or do not meet regional specifications for freedom from defect primarily because of roughness or poor form, and (b) all live trees of noncommercial species.

Roundwood products.--Logs, bolts, or other round sections (including chips from roundwood) cut from trees for industrial or consumer uses. (Note: Includes saw logs, veneer logs and bolts; cooperage logs and bolts; pulpwood; fuelwood; piling; poles; posts; hewn ties; mine timbers; and various other round, split, or hewn products.)

Salvable dead trees.--Standing or down dead trees that are considered merchantable by regional standards.

Saplings.--Live trees 1- to 5 inches d.b.h.

Sapling-seedling stands.--(See stand-size class.)

Saw log.--A log meeting minimum standards of diameter, length, and defect, including logs at least 8 feet long, sound and straight and with a minimum diameter outside bark (d.o.b.) for softwoods of 7 inches (9 inches for hardwoods) or other combinations of size and defect specified by regional standards.

Saw log portion.--That part of the bole of sawtimber trees between the stump and the saw log top.

Saw log top.--The point on the bole of sawtimber trees above which a saw log cannot be produced. The minimum saw log top is 7 inches d.o.b. for softwoods and 9 inches d.o.b. for hardwoods.

Sawtimber stands.--(See stand-size class.)

Sawtimber trees.--Growing-stock trees of commercial species containing at least a 12-foot saw log or two noncontiguous saw logs 8 feet or longer, and meeting regional specifications for freedom from defect. Softwoods must be at least 9 inches d.b.h. Hardwoods must be at least 11 inches d.b.h.

Sawtimber volume.--Net volume of the saw log portion of live sawtimber in board feet, International 1/4-inch rule, from stump to a minimum 7 inches top diameter outside bark (d.o.b.) for softwoods and a minimum 9 inches top d.o.b. for hardwoods.

Seedlings.--Live trees less than 1 inch d.b.h. that are expected to survive. Only softwood seedlings more than 6 inches tall and hardwood seedlings more than 1 foot tall are counted.

Short-log (rough tree).--Sawtimber-size trees of commercial species that contain at least one merchantable 8- to 11-foot saw log but not a 12-foot saw log.

Shrub biomass.--The total aboveground weight (including the bark) of selected shrubs and trees less than 1 inch d.b.h.

Site class.--A classification of forest land in terms of inherent capacity to grow crops of industrial wood based on fully stocked natural stands.

Site index.--An expression of forest site quality based on the total height of free-growing dominant or codominant trees of a representative species in the forest type at age 50.

Softwoods.--Coniferous trees, usually evergreen, having needles or scale-like leaves.

Stand.--A growth of trees on a minimum of 1 acre

of forest land that is stocked by forest trees of any size.

Stand-age class.--Age of the main stand. Main stand refers to trees of the dominant forest type and stand-size class.

Stand-area class.--The extent of a continuous forested area of the same forest type, stand-size class, and stand-density class.

Stand-size class.--A classification of forest land based on the size class of growing-stock trees on the area; that is, sawtimber, poletimber, or seedlings and saplings.

a. Sawtimber stands.--Stands at least 16.7 percent stocked with growing-stock trees, with half or more of total stocking in sawtimber or poletimber trees, and with sawtimber stocking at least equal to poletimber stocking.

b. Poletimber stands.--Stands at least 16.7 percent stocked with growing-stock trees of which half or more of this stocking is in poletimber and/or sawtimber trees, and with poletimber stocking exceeding that of sawtimber.

c. Sapling-seedling stands.--Stands at least 16.7 percent stocked with growing-stock trees of which more than half of the stocking is saplings and/or seedlings.

d. Nonstocked stands.--Stands in which stocking of growing-stock trees is less than 16.7 percent.

State land.--Land owned by States, or land leased to these governmental units for 50 years or more.

Stocking.--The degree of occupancy of land by trees, measured by basal area and/or the number of trees in a stand by size or age and spacing, compared to the basal area and/or number of trees required to fully utilize the growth potential of the land; that is, the stocking standard. A stocking percent of 100 indicates full utilization of the site and is equivalent to 80 square feet of basal area per acre in trees 5 inches d.b.h. and larger. In a stand of trees less than 5 inches d.b.h., a stocking percent of of 100 would indicate that the present number of trees is sufficient to produce 80 square feet of basal area per acre when the trees reach 5 inches d.b.h. Stands are grouped into the following stocking classes:

Overstocked stands.--Stands in which stocking of trees is 134.0 percent or more.

Fully stocked stands.--Stands in which stocking of trees is from 101.0 to 133.9 percent.

Medium stocked stands.--Stands in which stocking of trees is from 61.0 to 100.9 percent.

Poorly stocked stands.--Stands in which stocking of trees is from 16.7 to 60.9 percent.

Nonstocked areas.--Commercial forest land on which stocking of trees is less than 16.7 percent.

Timber removals from growing stock.--The volume of sound wood in growing-stock trees removed annually for forest products (including roundwood products and logging residues) and for other removals.

Timber removals from sawtimber.--The net board-foot volume of live sawtimber trees removed for forest products annually (including roundwood products and logging residues) and for other removals.

Timber products output.--All timber products cut from roundwood and byproducts of wood manufacturing plants. Roundwood products include logs, bolts, or other round sections cut from growing-stock trees, cull trees, salvable dead trees, trees on nonforest land, noncommercial species, sapling-size trees, and limbwood. Byproducts from primary manufacturing plants include slabs, edging, trimmings, miscuts, sawdust, shavings, veneer cores and clippings, and screenings of pulpmills that are used as pulpwood chips or other products.

Tree biomass.--The total aboveground weight (including the bark) of all trees from 1 to 5 inches in d.b.h., and the total aboveground weight (including the bark) from a 1-foot stump for trees more than 5 inches in diameter.

Tree size class.--A classification of trees based on diameter at breast height, including sawtimber trees, poletimber trees, saplings, and seedlings.

Unproductive forest land.--Forest land incapable of producing 20 cubic feet per acre of annual growth or of yielding crops of industrial wood under natural conditions because of adverse site conditions. (Note: Adverse conditions include shallow soils, dry climate, poor drainage, high elevation, steepness, and rockiness).

Upper stem portion.--That part of the bole of sawtimber trees above the saw log top to a minimum top diameter of 4 inches outside bark or to the point where the central stem breaks into limbs.

Urban and other areas.--Areas within the legal boundaries of cities and towns; suburban areas developed for residential, industrial, or recreational purposes; schoolyards, cemeteries, roads; railroads; airports; beaches; powerlines; and other rights-of-way; or other nonforest land not included in any other specified land use class.

Water.--(a) Bureau of the Census.--Permanent inland water surfaces, such as lakes, reservoirs, and ponds having 40 acres or more of area; streams, sloughs, estuaries, and canals one-eighth of a statute mile or more in width.

(b) Noncensus.--Permanent inland water surfaces, such as lakes, reservoirs, and ponds having 1 to 39.9 acres of area; streams, sloughs estuaries, and canals 120 feet to one-eighth of a statute mile wide.

Wooded pasture.--Improved pasture with more than 16.7 percent stocking in live trees but less than 25 percent stocking in growing-stock trees. Area is currently improved for grazing or there is other evidence of grazing.

Wooded strip.--An acre or more of natural continuous forest land that would otherwise meet survey standards for commercial forest land except that it is less than 120 feet wide.

BIBLIOGRAPHY

Belcher, David L.; Holdaway, Margaret R.; Brand, Gary J. A description of STEMS--the stand and tree evaluation and modeling system. Gen. Tech. Rep. NC-79. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1982. 18 p.

Hahn, Jerold T. Local net volume equations for Missouri. Gen. Tech. Rep. NC-15. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1975. 8 p.

Hahn, Jerold T. A local net volume equation for Iowa. Res. Note NC-199. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1976. 4 p.

Hahn, Jerold T. Tree volume and biomass equations for the Lake States. Res. Pap. NC-250. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1984. 10 p.

Robinson, Barry N.; Anderson, Gary D.; Cohen, Eli,; Gazdzik, Wally F.; Karpel, Lenny C.; Miller, Alan H.; Stein, Jack R. SIR scientific information retrieval users manual-version 2. Evanston, ILL: SIR Inc., 1980. p.

Smith, W. Brad and Weist, Carol A. A net volume equation for Indiana. Resour. Bull. NC-63. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1982. 7 p.

Hahn, Jerold T.; Hansen, Mark H.

Data bases for forest inventory in the North-Central Region. Gen. Tech. Rep. NC-101. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1985. 57 p.

Describes the data collected by the Forest Inventory and Analysis (FIA) Research Work Unit at the North Central Forest Experiment Station. Explains how interested parties may obtain information from the data bases either through direct access or by special requests to the FIA data base manager.

---

KEY WORDS: Forest survey, data retrieval, sampling procedures, data base management system.